




**City Of Fayetteville, Arkansas
2020 Woolsey Wet Prairie Sanctuary
Adaptive Management Strategy & Monitoring Report No. 14**



DECEMBER 2020




**CITY OF
FAYETTEVILLE
ARKANSAS**



**City of Fayetteville, Arkansas
2020 Woolsey Wet Prairie Sanctuary
Adaptive Management Strategy & Monitoring Report No. 14**



**Field Monitoring, Adaptive Management Guidance, & Report
Preparation Conducted by
Karen Willard, Bruce Shackleford, & Seth Pickens
Environmental Consulting Operations, Inc. Benton, Arkansas**



**"Much Appreciation" to Jeff Hickle of Jacobs
for his "hands on" hard work in making
adaptive management happen!**



"Special Thanks" to Dorothy Mangold for report cover photographs:

- Woolsey Wet Prairie Sanctuary sunset
- American bird grasshopper
- Blue grosbeak
- Blue-faced meadowhawk



DECEMBER 2020

TABLE OF CONTENTS

Section	Page
1.0 – Introduction & Project Background	1
1.1 – Individual Section 404 Permit No. 14207	1
1.2 – Mitigation Site Concept & Team	1
1.3 – Historical Rare Plant Species Observed at Woolsey Wet Prairie Sanctuary	3
1.4 – Historical Monitoring Strategies at Woolsey Wet Prairie Sanctuary	5
2.0 – 2020 Growing Season Observations	6
2.1 – Wetland Cell E-1 Observations	7
2.2 – Wetland Cell E-2 Observations	7
2.3 – Wetland Cell E-3 Observations	7
2.4 – Wetland Cell E-4 Observations	8
2.5 – Wetland Cell E-5 Observations	9
2.6 – Wetland Cell W-1 Observations	9
2.7 – Wetland Cell W-2 Observations	10
2.8 – Gas Line Easement Observations	11
2.9 – Overall Plant Species Richness Trends at Woolsey Wet Prairie	11
3.0 – 2020 Adaptive Management Activities	12
3.1 – Mowing & Hand Cutting/Pulling	12
3.2 – Prescribed Burning	12
3.3 – Hydrological Controls	13
3.4 – Historical Herbicide Applications	14
4.0 – Recommended Adaptive Management Activities for 2021	16
5.0 – 2021 Existing Surplus Wetland Mitigation Credits	17
5.1 – Background and Overview	17
6.0 – Academic Research and Guidance	18
8.0 – Appendices	19
Appendix I – 2020 Stop Log Data for Hydrological Controls in Wetland Cells	
Appendix II – Woolsey Wet Prairie Sanctuary Herbicide Mixing/Application Rates	
Appendix III – Tracking Species Aerial Images	
Appendix IV – Invasive Species Aerial Images	
Appendix V – Woolsey Wet Prairie 2020 Master Plant Species List	
Appendix VI – Woolsey Wet Prairie 2020 Adaptive Management Activities	
Appendix VII – 2020 Woolsey Wet Prairie Surplus Wetland Credit Ledger Report	
List of Figures	Page
Figure 1. Woolsey Wet Prairie Aerial Photograph	2
List of Tables	Page
Table 1 – Typical Seasonal Stop Log Settings	13
Table 2 – 2021 Woolsey Wet Prairie Adaptive Management Tentative Schedule	16

CITY OF FAYETTEVILLE, ARKANSAS
WOOLSEY WET PRAIRIE SANCTUARY
ADAPTIVE MANAGEMENT STRATEGY & MONITORING REPORT NO. 14

1.0 – INTRODUCTION & PROJECT BACKGROUND

The City of Fayetteville, Arkansas' Wastewater System Improvement Project (WSIP) entailed improvements to the City's sewer collection system, upgrading the Paul Noland Wastewater Treatment Plant (WWTP), and construction of the new Westside WWTP. Construction activities for these improvements occurred during a period from 2005 through 2010. The WSIP primary purpose was to implement corrective actions to eliminate/reduce odor and overflow problems associated with the Noland Plant and collection system and to provide wastewater treatment to areas outside the treatment area while reducing the total hydraulic loading to the system. WSIP activities involved discharges of fill into "Waters of the U.S." within the Illinois River Watershed and the Beaver Reservoir Watershed (within the White River Basin); therefore, U.S. Army Corps of Engineers (COE) permitting under Section 404 of the Clean Water Act (CWA) was required.

1.1 – Individual Section 404 Permit No. 14207

On March 10, 2005, the City of Fayetteville received Individual Section 404 Permit No. 14207 from the COE Little Rock District for the portion of the WSIP in the Illinois River Watershed (west side) that involved 36 stream crossings and 15 wetland crossings during construction of the new Westside WWTP, sewer lines, and improvements to Broyles Road. The 404 permit required wetland compensatory mitigation due to the permanent alteration of 8.87 acres of wetlands. As required by the COE, the wetland mitigation site was deed restricted in perpetuity to guarantee preservation of the wetlands and upland buffers. A certified copy of the Notice of Deed Restriction was recorded with the Washington County Registrar of Deeds on January 5, 2007. The City of Fayetteville is required to manage and maintain the property as a compensatory wetland mitigation site in perpetuity.

1.2 – Mitigation Site Concept and Team

The 43.65-acre wetland mitigation site (Figure 1) is located immediately to the north of the Westside WWTP. McGoodwin, Williams, and Yates Consulting Engineers, Inc. (MWY) provided civil engineering design of hydrological features and Environmental Consulting Operations, Inc. (ECO, Inc.) provided ecological feature design, site management, and monitoring. Brasfield and Gorrie General Contractors completed construction of earthen berms and water level control structures. Operation of hydrological controls, herbicide applications, fire line installations, and mowing are managed through Jacobs, who also maintains the City's wastewater utility system. Prescribed burns are contracted by the City through an informal bidding process. ECO, Inc. oversees Section 404 permit compliance, develops prescribed burn and herbicide application specifications, identifies problem areas where invasive plant species need to be controlled, and conducts annual monitoring and site adaptive management strategy development.

Modifications to the existing hydrology at the mitigation site were achieved via the construction of low elevation perimeter earthen berms designed to provide a mechanism for water retention within seven wetland cells. The micro-topography within the wetland cells was not disturbed or altered during the construction of the berms. The presence of depressions between prairie mounds provides a diversity of hydrology, which, in turn provides diversity in wetland and upland plant communities.

Figure 1 – Woolsey Wet Prairie Aerial Photograph

Water level control structures with stop logs were constructed within the berms in order to provide the ability to hold and to release water as needed in order to maintain inundation/saturation within the wetland cells for the purpose of optimizing wetland plant community diversity. Construction of the earthen berms resulted in two cells (W-1 and W-2) within the West Mitigation Site, and five cells (E-1 through E-5) within the East Mitigation Site. The west and east mitigation sites are separated by a gas pipeline easement that is 60 feet in width. The easement has undergone the same adaptive management as the remaining acreage on the deed restricted property. A perimeter buffer zone has been established around the perimeter of the deed-restricted mitigation site where herbicide applications and prescribed burns are performed to prevent migration of non-native invasive plant species into the mitigation area. Therefore, a total of approximately 65-acres are managed on the property.

In 2006, Bruce Shackelford, ECO, Inc. President, named the mitigation site “Woolsey Wet Prairie Sanctuary” in honor of Samuel Gilbert Woolsey, whose family settled the property in 1830 when prairies were abundant in Northwest Arkansas.

The Woolsey Wet Prairie Sanctuary is part of the original prairie of Prairie Township, Fayetteville, Arkansas that extended all the way to the Prairie Grove and Lincoln areas in Washington County. Conversion of an estimated 100,000 acres of both wetland and upland prairie habitat in Northwest Arkansas to production of wheat in the late 1800’s and early 1900’s was the beginning of the decimation of prairie habitat, America’s most endangered ecosystem. Fire suppression and the introduction of non-native plant species have also contributed greatly to the near complete extirpation of prairie habitat.

1.3 – Historical Rare Plant Species Observed at Woolsey Wet Prairie Sanctuary

Eleven plant species tracked as elements of conservation concern (rare species) by the Arkansas Natural Heritage Commission (ANHC), have been found to naturally occur at the wetland mitigation site. These rare plants include sedges (family *Cyperaceae*), milkwort (family *Polygalaceae*), Mexican white sage (family *Asteraceae*), and Hawthorn (genus *Crataegus*), and are characteristic of unplowed tall grass wet prairie remnants.

***Carex aggregata* (cluster sedge) – G5S1** – This sedge is known in Arkansas only from a few sites in Benton, Carroll, Fulton, Newton, Sharp, and Washington counties. It typically grows in low open woodlands or seasonally wet grasslands. At Woolsey Wet Prairie, it is scattered in seasonally wet areas that are not inundated for long periods. It has been historically and presently observed in all seven wetland cells and within the gas line easement at Woolsey Wet Prairie.

***Polygala incarnata* (pink milkwort) – G5S1S2** – This rare species of wildflower is known in Arkansas from remnant prairies and other historically open grassland habitats, such as glades and savannas. It is known from scattered counties in Arkansas, but most of the records of many of the sites where it was historically found have since been destroyed. A single plant was found on a prairie pimple mound in wetland cell E-4 in 2012 and was not observed in 2013. In 2014, this population increased to six plants, but none were observed from 2014 through 2017. It is thought that this plant's absence was due to shading out associated with the increase in woody plant densities observed during 2015 and 2016. Timber mulching activities were conducted during 2017 to reduce the over shading effect of woody vegetation. During 2018, a single plant was observed for the first time in the buffer zone south of wetland cell W-2. However, during 2019 and 2020, no individuals of this species were observed at the site. It has not been ascertained whether or not this species still exists at the site.

***Carex scoparia* var. *scoparia* (pointed sedge) – G5S1S2** – This species is very rare in Arkansas and is known only from prairie-associated wetlands in Washington and Benton counties. Historically, this plant was observed as a single clump that has persisted in wetland cell W-1. During 2018 and 2019, it was observed in wetland cells E-4 and E-5, W-1 and W-2. In 2020, it persists in wetland cells E-5, W-1 and W-2, but was not observed in E-4.

***Carex arkansana* (Arkansas sedge) – G4S2** – This uncommon sedge is known in Arkansas from wet prairie remnants, open hydric oak flatwoods, and similar open wetland habitats (ANHC, 2014). While it has no wetland indicator status code in the United States Department of Agriculture (USDA) Plants Database, it is listed by Yatskievych (1999) as occurring primarily in bottomland prairies and moist depressions of upland prairies. At Woolsey Wet Prairie, it is scattered in seasonally wet areas that are only inundated for short periods. Historically, this plant species has been observed in all seven wetland cells and the gas line easement, but was absent during 2018 for the first time in monitoring history from wetland cell E-3. Monitoring activities during 2019 produced similar results, with the species being observed in all wetland cells and the gas line easement, with the exception of wetland cell E-3. Monitoring activities during 2020 revealed its presence in all wetland cells.

***Carex opaca* (opaque prairie sedge) – G4S2S3** – This rare sedge is primarily associated with unplowed, wet tall grass prairie remnants in Arkansas (ANHC, 2014). While it has no wetland indicator status code in the USDA Plants Database, it is listed by Yatskievych (1999) as primarily occurring in bottomland prairies, moist depressions of upland prairies, and margins of fens. At Woolsey Wet Prairie, it is scattered in seasonally wet areas that are not inundated for long periods. Historically and presently, it has consistently been found in all seven wetland cells and the gas line easement at Woolsey Wet Prairie through monitoring year 2020.

***Carex fissa* var. *fissa* (hammock sedge) – G4S1** – Prior to its discovery at Woolsey Wet Prairie, this rare sedge was known in Arkansas from only two sites in Franklin and Lonoke Counties, where it occurs in prairie-associated wetlands (ANHC, 2014). At Woolsey Wet Prairie, it has historically been found in three naturally occurring prairie swales in wetland cells E-2, W-1, and W-2. During 2015, it was absent from wetland cell E-2, but returned during 2016 and was observed again during 2018. During 2015 and 2016, it was absent from wetland cell W-1, but returned and was observed again in 2018. It is possible that this plant disappeared due to shading associated with the increase in woody plant densities observed during 2015 and 2016, and the timber mulching activities conducted during 2017 aided in this species return to the site. In 2018, it was observed for the first time within the gas line easement. This species was observed in 2019 in wetland cells W-1, W-2, and within the gas line easement, but was not observed in wetland cell E-2. It was observed in wetland cells W-1 and W-2 during 2020, but was not observed in the gas line, or wetland cell E-2.

***Carex pellita* (woolly sedge) – G5S1S2** – Prior to its collection at Woolsey Wet Prairie, this species was known to be extant at a single Arkansas locality, in a fen in Marion County. It has since been found at three other sites in Benton, Washington, and Marion counties. At Woolsey Wet Prairie, it has been historically observed at seasonally wet areas in wetland cells E-4, E-5, W-1, and W-2. It was absent from wetland cell W-1 in 2018. During 2019, this species was observed in wetland cells E-4, E-5, W-1, W-2, and within the gas line easement for the first time, but was absent from the gas line in 2020.

***Eleocharis wolfii* (Wolf's spikerush) – G3G4S3** – This wetland sedge occurs in Arkansas primarily in wet areas in unplowed tall grass prairie remnants, but can persist in wet, open areas in landscapes that were formerly dominated by prairie vegetation (ANHC, 2014). At Woolsey Wet Prairie, this species has historically been found in several naturally occurring swales within wetland cells E-2, E-3, E-4, E-5, W-1, and W-2. It was not observed in wetland cell E-5 during 2015 and 2016, possibly due to shading associated with the increase in woody plant densities observed during 2015 and 2016, prior to the timber mulching activities conducted during 2017. During 2019 and 2020, this species was observed in wetland cells E-2, E-3, E-4, E-5, W-1, W-2, and within the gas line easement.

***Rhynchospora macrostachya* (tall horned beaksedge) – G4S2** – Prior to its collection at Woolsey Wet Prairie, this species was known in Arkansas only from a few scattered historical collections from remnant prairies. It has since also been found in several prairie-associated wetlands in Franklin County. At Woolsey Wet Prairie, it was first known from two natural prairie swales prior to construction of the berms. In the fall of 2006, ECO, Inc. gathered seeds and successfully propagated over 50 specimens during the 2007 growing season that were transplanted into marsh areas at the site during 2008. A 90 percent survival rate was observed and transplanted specimens produced large seed heads by the end of the 2008 growing season. Historically, this species has been observed in wetland cells E-2, E-4, E-5, W-1, W-2, and within the gas line easement. In 2015 and 2018, it was absent from wetland cell E-2, and in 2018 it was absent from wetland cells E-4 and E-5. During 2019 and 2020, this species was observed in wetland cells E-5, W-1, W-2, and within the gas line easement, but still absent from wetland cell E-2.

***Crataegus reverchonii* (Reverchon’s hawthorn) – G4S1** – This small tree has been confirmed to occur in Arkansas only in Benton and Washington counties at low prairies or woodlands. It is primarily a western species. Specimens at Woolsey Wet Prairie appear to be *Crataegus reverchonii* subsp. *palmeri*, but both *palmeri* and the subspecies *reverchonii* have been reported for northwestern Arkansas. Additional study is needed to determine if both subspecies are present at the site. Historically, it has been observed within wetland cells W-1, E-4, E-5, and W-2. During 2019, this species was only observed within wetland cell W-1. However, it was observed in wetland cells E-4, W-1, and W-2 during the 2020-monitoring year.

***Artemisia ludoviciana* var. *mexicana* (Mexican white sage) – G5T5S1S2** – This species is known to occur in Arkansas in dry grasslands and glades in a few counties in the northwestern part of the state. It was last documented from the Fayetteville area in 1954, when it was collected from “West Mountain” (a site believed to be about two miles east of Woolsey Wet Prairie). Historically, this species was observed as two distinct patches located in a fencerow and field margin along the south side of Persimmon Street, just west of Owl Creek. During 2018, Jeff Hickle of Jacobs collected and propagated seeds from these individuals and successfully transplanted them within the boundaries of the site during early 2019 in the buffer areas north of wetland cells E-1, E-2, west of W-1, and east of E-3. During 2020 individual plant species were observed in the buffer areas of wetland cells W-1 and W-2.

Key to ANHC Species Category Rankings:

G3 – Either very rare and local throughout its range or found locally (even abundantly at some of its locations) in a restricted range

G4 – Apparently secure globally, though it may be quite rare in parts of its range, especially at the periphery

G5 – Demonstrably secure globally, although it may be quite rare in parts of its range, especially at the periphery

T – Subspecies or variety rank (ex. G5T4 applies to a subspecies with a global species rank of G5, but with a subspecies rank of G4)

S1 – Critically imperiled because of extreme rarity or because of some factor(s) making it especially vulnerable to extirpation

S2 – Imperiled because of rarity (6 to 20 known extant populations) or because of some factor(s) making it especially vulnerable to extirpation

S3 – Rare throughout the state or found locally in a restricted region of the state, or because of other factors making it vulnerable to extirpation

1.4 – Historical Monitoring Strategies at Woolsey Wet Prairie Sanctuary

As part of the terms and conditions included in the City of Fayetteville’s Section 404 permit, seven annual reports on the status of the mitigation site were required to be submitted to the COE. The first annual wetland monitoring report was due December 31st after the first growing year, and each year thereafter, for a total of seven years. The first monitoring year was 2007 and the seventh annual monitoring report was completed in December 2013. Initially, the COE required intensive monitoring activities at 47 monitoring stations for the first seven years. Since that time, ECO, Inc. has

transitioned to an abbreviated methodology strategy that focuses more on where adaptive management activities are needed on a cell-by-cell basis in lieu of the 47 monitoring stations. This serves to assure that the City of Fayetteville continues to meet COE Section 404 permit required ecological performance standards and maintains eligibility to use surplus wetland credits for city infrastructure improvements that permanently alter wetlands.

ECO, Inc. conducted periodic site visits throughout the 2020-growing season to observe and evaluate the effectiveness of herbicide applications for control of invasive plant species, to evaluate plant succession, to evaluate performance standards status, and to observe the status of site hydrology. ECO, Inc. contracted Karen Willard from the University of Arkansas herbarium to conduct vegetation monitoring events during June and September to document the presence, densities, and locations of stands of non-native/invasive plant species and rare plant species tracked as elements of conservation concern by the ANHC. Karen marked Google Earth generated aerial photographs in the field to show locations of non-native/invasive and rare plant species. The non-native/invasive species aerial maps were used by Jacobs during the growing season as a guide for areas most heavily targeted for herbicide applications. The rare species aerial maps were used during the academic research project guidance, mitigation, and approval process.

The results of the 2020 field observations and monitoring data are compiled herein for the purpose of evaluating the degree of success in controlling non-native/invasive plant species that threaten the rare prairie ecosystem at Woolsey Wet Prairie and to develop and prioritize an action item list for adaptive management activities and goals for the upcoming 2021 growing season.

In prior years, monitoring activities focused largely on evaluating species richness within each wetland cell and for the overall site. During the 2018, 2019, and 2020 growing seasons, an increased focus was placed upon the evaluation of adverse impacts to the native prairie plants caused by persistent invasive plant species and the woody plant succession that was adversely affecting the prairie character of the site. Observations made during the 2017 and 2018 monitoring activities were used to develop a more aggressive plan of action for controlling invasive and woody plant species in 2018, 2019, and 2020. Positive results were achieved via this approach that will be continued throughout the 2021-growing season.

2.0 – 2020 GROWING SEASON OBSERVATIONS

The following sections describe observations for each wetland cell during the 2020 growing season. For purposes of adaptive management at Woolsey Wet Prairie, invasive plant species include both native and non-native plants that have the potential to outcompete native prairie and wetland plant species in a manner and degree that poses an ecological threat to sustaining the wetland prairie ecosystem.

2.1 – Wetland Cell E-1 Observations

E-1 Rare Species

Three rare plant species, cluster sedge, Arkansas sedge, and opaque prairie sedge were observed in wetland cell E-1. The locations of these rare plant species are shown on the aerial photograph contained within Appendix III.

E-1 Invasive Species

Fourteen invasive species were observed in this wetland cell in 2020 that need continued management; however, the densities of these invasives have been greatly reduced as compared to previous years. In particular, small carpetgrass (*Arthraxon hispidus*), Bermuda grass (*Cynodon dactylon*), Queen Anne's lace (*Daucus carota*), southern crabgrass (*Digitaria ciliaris*), smooth crabgrass (*Digitaria ischaemum*), Korean bushclover (*Kummerowia stipulacea*), Japanese Bushclover (*Kummerowia striata*), sericea lespedeza (*Lespedeza cuneata*), white sweet clover (*Melilotus albus*), Callery pear (*Pyrus calleryana*), curly dock (*Rumex crispus*), bitter dock (*Rumex obtusifolius*), tall fescue (*Schedonorus arundinaceus*), and Johnsongrass (*Sorghum halepense*) are all persisting. The locations of the non-native/invasive plant species are marked on the E-1 Aerial Photograph contained within Appendix IV. The management team has experienced significant success in 2020 in terms of reducing density and species richness of invasive plant species.

2.2 – Wetland Cell E-2 Observations

E-2 Rare Species

Four rare plant species, cluster sedge, Arkansas sedge, opaque prairie sedge, and Wolf's spikerush were observed in wetland cell E-2. In 2017, tall horned beaksedge was observed and in 2018 hammock sedge was observed; however, neither of these rare plant species were observed as being present during 2019 or 2020. The locations of these rare plant species are shown on the aerial photograph contained within Appendix III.

E-2 Invasive Species

Eighteen invasive species were observed in this wetland cell in 2020 that need continued management; however, the densities of these invasives have been greatly reduced as compared to previous years. In particular, small carpetgrass, hairy brome (*Bromus racemosus*), Bermuda grass, Queen Anne's lace, smooth crabgrass, Korean bushclover, sericea lespedeza, Japanese honeysuckle (*Lonicera japonica*), white sweet clover, callery pear, multiflora rose (*Rosa multiflora*), Himalayan blackberry (*Rubus serissimus*), curly dock, bitter dock, tall fescue, Johnsongrass, spiny Sowthistle (*Sonchus asper*), and narrowleaf cattail (*Typha angustifolia*) are all persisting. The locations of the non-native/invasive plant species are marked on the E-2 Aerial Photograph contained within Appendix IV. The management team has experienced significant success in 2020 in terms of reducing density and species richness of invasive plant species.

2.3 – Wetland Cell E-3 Observations

E-3 Rare Species

Four rare plant species, cluster sedge, Arkansas sedge, opaque prairie sedge, and Wolf's spikerush were observed in wetland cell E-3. Arkansas sedge, which has historically been observed within this

cell, was not observed in 2018 or 2019, but was found to once again be present during 2020. The locations of these rare plant species are shown on the aerial photograph contained within Appendix III.

E-3 Invasive Species

Fifteen invasive species were observed in this wetland cell in 2020 that need continued management; however, the densities of these invasives have been greatly reduced as compared to previous years. In particular, small carpetgrass, hairy brome, poverty brome (*Bromus sterilis*), nodding thistle (*Carduus nutans*), common thistle (*Cirsium vulgare*), Bermuda grass, Queen Anne's lace, southern crabgrass, Korean bushclover, Japanese bushclover, sericea lespedeza, white sweet clover, Callery pear, curly dock, and tall fescue are all persisting. The locations of the non-native/invasive plant species are marked on the E-3 Aerial Photograph contained within Appendix IV. The management team has experienced significant success in 2020 in terms of reducing density and species richness of invasive plant species.

2.4 – Wetland Cell E-4 Observations

E-4 Rare Species

Six rare plant species were observed in wetland cell E-4. Cluster sedge, Arkansas sedge, and opaque prairie sedge are uncommon in this cell and are scattered in low areas that are not inundated for long durations. A single colony of woolly sedge observed in this cell in 2013 has persisted and expanded. In 2018, pointed sedge was observed for the first time within this cell and was observed to have persisted in 2019; however, it was not observed during 2020. Wolf's spikerush occurs scattered at the edge of open marshes. Pink milkwort was observed on a pimple mound in 2012 and 2014, but was not observed in 2013 or 2015 - 2020. However, this type of population fluctuation is not uncommon in annual species. A single plant of Reverchon's hawthorn, which has historically been observed near the west side of the cell since 2014 was not observed during 2019, but was once again observed during 2020. Tall horned beaksedge and hammock sedge, which have been historically observed within this cell, were found to not be present in 2018, 2019, or 2020. The locations of these rare plant species are shown on the aerial photograph contained within Appendix III.

E-4 Invasive Species

Twenty invasive species were observed in this wetland cell in 2020 that need continued management; however, the densities of these invasives have been greatly reduced as compared to previous years. In particular, small carpetgrass, hairy brome, poverty brome, nodding thistle, common thistle, Bermuda grass, Queen Anne's lace, Korean bushclover, Japanese bushclover, sericea lespedeza, Japanese honeysuckle (*Lonicera japonica*), white sweet clover, Japanese stiltgrass (*Microstegium vimineum*), Callery pear, multiflora rose, Himalayan blackberry, curly dock, bitter dock, tall fescue, and spiny sowthistle are all persisting. The locations of the non-native/invasive plant species are marked on the E-4 Aerial Photograph contained within Appendix IV.

This wetland cell has approximately 0.54 acres of designated Tree Preservation Areas where woody vegetation is allowed to grow in order to meet COE compensatory requirements for permanent alterations to forested wetlands during construction of the WSIP. Otherwise, woody vegetation is controlled at other locations to prevent a prairie ecosystem from being converted into a forested ecosystem.

2.5 – Wetland Cell E-5 Observations

E-5 Rare Species

Seven rare plant species were observed in wetland cell E-5. Cluster sedge is rare and scattered on the east side of the cell. Arkansas sedge and opaque prairie sedge are uncommon and are scattered in low areas that are not inundated for long durations. A single colony of woolly sedge observed in 2013 has persisted and expanded in the northeast portion of the cell. Wolf's spikerush occurs scattered at the edge of open marshes on the south and east portions of the cell. Pointed sedge was observed for the first time within this cell in 2018 and was found to have persisted during 2019 and 2020. Tall horned beaksedge, which has been historically observed within this cell, but was not observed during 2018, was once again observed during 2019 and 2020. The locations of these rare plant species are shown on the aerial photograph contained within Appendix III.

E-5 Invasive Species

Fourteen invasive species were observed in this wetland cell in 2020 that need continued management; however, the densities of these invasives have been greatly reduced as compared to previous years. In particular, small carpetgrass, hairy brome, nodding thistle, common thistle, Bermuda grass, Queen Anne's lace, southern crabgrass, Japanese bushclover, sericea lespedeza, Japanese honeysuckle, curly dock, tall fescue, spiny sowthistle, and narrowleaf cattail are all persisting. The locations of the non-native/invasive plant species are marked on the E-5 Aerial Photograph contained within Appendix IV.

This wetland cell has approximately 0.75 acres of designated Tree Preservation Areas where woody vegetation is allowed to grow in order to meet COE compensatory requirements for permanent alterations to forested wetlands during construction of the WSIP. Otherwise, woody vegetation is controlled at other locations to prevent a prairie ecosystem from being converted into a forested ecosystem.

2.6 – Wetland Cell W-1 Observations

W-1 Rare Species

Seven rare plant species were observed in wetland cell W-1. The single clump of pointed sedge first observed in this cell in 2012 is still persisting and has expanded. Cluster sedge, Arkansas sedge, and opaque prairie sedge are uncommon in this cell and are scattered in low areas that are not flooded for long durations. Tall horned beaksedge and Wolf's spikerush occur scattered at the edge of open marshes. Reverchon's hawthorn was found to persist outside the berm on the west side of this cell. A few individuals of hammock sedge that in previous years had been shaded out by willows were observed to be present in 2018 following the 2017 timber mulching activities. These individuals were observed again during 2019 and 2020. Woolly sedge, which has been historically observed within this cell, was found to not be present in 2018, 2019, or 2020. The locations of these rare plant species are shown on the aerial photograph contained within Appendix III.

W-1 Invasive Species

Eighteen invasive species were observed in this wetland cell in 2020 that need continued management; however, the densities of these invasives have been greatly reduced as compared to previous years. In particular, small carpetgrass, hairy brome, nodding thistle, Bermuda grass, Queen

Queen Anne's lace, southern crabgrass, Korean bushclover, Japanese bushclover, sericea lespedeza, Japanese honeysuckle, white sweetclover, callery pear, Himalayan blackberry, curly dock, bitter dock, tall fescue, spiny sowthistle, and narrow-leaf cattail are all persisting. The locations of the non-native/invasive plant species are marked on the W-1 Aerial Photograph contained within Appendix IV.

This wetland cell has approximately 1.2 acres of designated Tree Preservation Areas where woody vegetation is allowed to grow in order to meet COE compensatory requirements for permanent alterations to forested wetlands during construction of the WSIP. Otherwise, woody vegetation is controlled at other locations to prevent a prairie ecosystem from being converted into a forested ecosystem.

2.7 – Wetland Cell W-2 Observations

W-2 Rare Species

Eight rare plant species occur in wetland cell W-2, with this cell being the most rich in rare plant species. Several clumps of cluster sedge were found in the central portion of the cell. Arkansas sedge, which in 2019 was only observed as a few individuals, was common throughout the cell. Opaque prairie sedge is fairly common and is scattered throughout the cell in low areas that are not inundated for long durations. Historically, a large population of hammock sedge existed in the southern half of this cell, but during 2018, 2019, and 2020 only a few individuals were observed. Several colonies of woolly sedge occur in the northeast, southwest, and southeast portions of this cell. Wolf's spikerush is fairly common and is scattered throughout the cell. Pointed sedge was observed for the first time within this cell in 2018 as scattered individuals. During 2019 and again in 2020 it was observed just outside the berm on the northwest corner and inside the berm in the northeast portion of the cell. In 2018, Pink milkwort was observed for the first time outside of the southern berm, but was not observed during 2019 or 2020. A single large, fruiting tree of Reverchon's hawthorn was observed on the west side of the wooded area outside the berm on the west edge of this cell. Tall horned beaksedge was observed in the southwest portion of the cell. The locations of these rare plant species are shown on the aerial photograph contained within Appendix III.

W-2 Invasive Species

Nineteen invasive species were observed in this wetland cell in 2020 that need continued management; however, the densities of these invasives have been greatly reduced as compared to previous years. In particular, small carpetgrass, poverty brome, nodding thistle, Bermuda grass, Queen Anne's lace, southern crabgrass, Korean bushclover, Japanese bushclover, sericea lespedeza, Japanese honeysuckle, white sweetclover, Callery pear, Himalayan blackberry, curly dock, bitter dock, tall fescue, spiny sowthistle, and narrow-leaf cattail are all persisting.

The wooded area outside the berm on the west edge of this cell has historically been very thick with invasive plants including multiflora rose, Chinese privet (*Ligustrum sinense*), bush honeysuckle (*Lonicera maackii*), Himalayan blackberry, Japanese honeysuckle, and winter-creeper (*Euonymus fortunei*). These species were not observed in this area during 2020. In October 2017, the very invasive Chinese Silver Grass (*Miscanthus sinensis*) was found for the first time at the site in the southeast corner of Wetland Cell W-2 just west of the old farm pond. The tall clumps of Chinese Silver Grass were treated using the herbicide Outrider. Following the spring 2018 burn, individuals of

this plant reemerged from dormancy and were again treated with very positive effects. During 2019 and 2020, no individuals of this species were observed. The locations of the non-native/invasive plant species are marked on the W-2 Aerial Photograph contained within Appendix IV.

2.8 – Gas Line Easement Observations

Gas Line Easement Rare Species

A 60-foot wide high-pressure gas line easement separates the East Wetland Mitigation Cells from the West Wetland Mitigation cells. This easement is not included within the acreage designated for the deed-restricted wetland mitigation site. However, it is managed in the same manner as the wetland cells for the purpose of controlling invasive plant species and preserving the rare plant species.

Six rare plants occur within the gas line easement. Arkansas sedge, opaque prairie sedge, and cluster sedge are fairly common and are scattered in low areas that are not inundated for long durations. A large colony of woolly sedge occurs in the southern half of this cell. Tall horned beaksedge was observed as a few individuals in the central portion of the cell. Wolf's spikerush occurs scattered throughout the cell at the edges of open marshes. Hammock sedge, which has been historically observed within this cell, was found to not be present in 2020. The locations of these rare plant species are shown on the aerial photograph contained within Appendix III.

Gas Line Easement Invasive Species

Eighteen invasive species were observed in this wetland cell in 2020 that need continued management; however, the densities of these invasives have been greatly reduced as compared to previous years. In particular, small carpetgrass, hairy brome, nodding thistle, Bermuda grass, Queen Anne's lace, southern crabgrass, smooth crabgrass, Korean bushclover, Japanese bushclover, sericea lespedeza, Japanese honeysuckle, Callery pear, multiflora rose, Himalayan blackberry, curly dock, bitter dock, tall fescue, and spiny sowthistle are all persisting. The locations of the non-native/invasive plant species are marked on the Gas Line Aerial Photograph contained within Appendix IV.

2.9 - Overall Plant Species Richness Trends at Woolsey Wet Prairie

Management, inventory, and monitoring work findings in 2020 resulted in the addition of seven new plant species not previously observed at the Woolsey Wet Prairie. Among the 2020 additions were Philadelphia fleabane (*Erigeron philadelphicus*), lanceleaf loosestrife (*Lysimachia lanceolata*), Japanese stiltgrass (*Microstegium vimineum*), beefsteak plant (*Perilla frutescens*), soft goldenaster (*Bradburia pilosa*), sumpweed (*Iva annua*), and splitbeard bluestem (*Andropogon ternarius Michx.*).

To date, a total of 496 plant species have been observed at Woolsey Wet Prairie since 2001. Of these, 99 (20%) species are considered to be non-native and/or invasive species, and 395 (80%) are considered to be native plant species. Because different species emerge at different times of the year, and the fact that some plant species may not emerge every year, this does not mean that 496 plant species can be observed at any one moment in time. However, new plant species are appearing each year and a 1,055 percent increase in plant species richness has been observed since 2005 when only 47 plant species were observed at the site. The 2020 Woolsey Wet Prairie Master Plant Species List is contained within Appendix V.

3.0 – 2020 ADAPTIVE MANAGEMENT ACTIVITIES

ECO, Inc. develops annual adaptive management goals and objectives and works as a team with Jacobs who is the party that provides boots on the ground services at Woolsey Wet Prairie to implement adaptive management techniques such as mowing, herbicide application, and hand cutting/pulling. A comprehensive summary of the adaptive management activities completed at Woolsey Wet Prairie during 2020 is contained within Appendix VI.

3.1 – Mowing & Hand Cutting/Pulling

The mowing at the site is aimed toward invasive species such as tall fescue, Johnsongrass, Dallis grass (*Paspalum dilatatum*), Queen Anne's lace, ragweed (*Ambrosia spp.*), and sericea lespedeza. When necessary, stands of these species are mowed to a height of 10-12 inches as they begin to mature, but before they form seed heads. This is intended to prevent the dispersal of additional seeds into the site seed bank from invasive species. As a precautionary measure, pastures to the north and west of the mitigation site are mowed to reduce the risk of seed dispersal of undesirable plant species from unrestored areas. The trails on top of the earthen berms are routinely mowed for the purpose of maintaining site access by the public and to control invasive plant species from forming seeds.

3.2 – Prescribed Burning

To date, twelve prescribed burns have been conducted at Woolsey Wet Prairie on the following dates, February 29, 2008, February 19, 2009, December 16, 2009, March 18, 2011, March 13, 2012, March 3, 2013, March 13, 2014, March 23, 2015, March 5, 2016, February 24, 2017, March 8, 2018, March 18, 2019, and February 21, 2020. ECO, Inc. routinely develops prescribed burn specifications and a prescribed burn contractor is selected through an informal bid process.

For ecological restoration, fire has become recognized as a valuable vegetation management tool that can be used to enhance community diversity. Fire removes much of the surface layer of decaying vegetation "thatch" that covers the ground. Many native plant species require sunlight to germinate, while others actually require fire to germinate. Prescribed burns aide in preventing woody plant encroachment and maintains the wet prairie habitat, depending upon the time of year of the burn, and the site hydrology at the time of the burn. The volunteer woody plant growth has primarily occurred in the wetter areas where inundation, and/or moist soils protect woody plants from fire.

With the objective of encouraging Native Warm Season Grasses (NWSG) and suppressing hardwood sprouts, the most effective burn period at Woolsey Wet Prairie has been found to be during the February to March time period. Ideally, this will occur during the transition from the Late Dormant to Dormancy Break periods. At that time, most of the warm season species will still be dormant and there will be adequate fuel from the vegetation killed by prior winter cold weather.

Prairies existed for thousands of years, primarily because Native Americans burned the local landscapes. Research has shown that they sometimes chose the most severe weather to ignite fires in order to maintain open grasslands that attracted the large herbivores they hunted. Some fires were done during drought conditions that greatly inhibited the plant succession, killing the majority of the trees in the fire's path. What remained were open grassland prairies with scattered post oak (*Quercus stellata*) savannas (very low density tree areas) that had an understory of native prairie grasses and forbs. The post oaks prevailed because they were the most fire-resistant tree species in the Ozarks.

Unlike the use of fire by Native Americans during drought conditions, the prescribed burns at Woolsey Wet Prairie must be conducted under safe conditions due to the urban-interface setting, and when there are no burn bans issued by the Arkansas Forestry Commission.

Prescribed burns help maintain a fire dependent prairie ecosystem, but they are likely not as hot and as intense as historical fires of Native American origin and do not kill a large number of trees. Furthermore, because Woolsey has an abundance of wetland areas, thorough burning of stands of black willow (*Salix nigra*) and green ash (*Fraxinus pennsylvanica*) are often difficult to achieve due to moisture levels in the ground and vegetation. Therefore, the prescribed burning is supplemented with herbicide applications to control the woody vegetation.

3.3 – Hydrological Controls

All wetland cells (with the exception of wetland cell W-2) have water level control structures. The structures have stop logs consisting of two dimensions; five-inch and seven-inch heights. This allows for control of water levels within the wetland cells within two-inch increments, depending upon the configuration of the stop logs and the amount of rainfall. In general, the stop log configurations are set to: 1) maintain surface water within portions of each wetland cell; 2) maintain non-inundated areas that have saturated surface soils; and 3) preclude overflows over the berms that would result in berm erosion. Maximum water retention within the wetland cells is not desirable, as it may create conditions not suitable for maintaining rare wetland sedge and rush species that cannot survive in periods of prolonged inundation.

Management of hydrology is an important tool in vegetation community diversity optimization because plant zonation occurs along water depth and soil saturation gradients. Consequently, variations in water depth and degree of soil saturation lead to variations in species composition. In summary, for management of hydrology, the major emphasis will be to recreate natural hydrological regimes in a manner to limit productivity of any single species from becoming excessively high, while at the same time, enriching biodiversity. The strategy for management of hydrology has not only included considerations for the volume of water retained, but also the time of the year water is retained. It is vital to retain water during the growing season in order to maintain areas of soil saturation and/or inundation to support desirable wetland vegetation. Table 1 below shows the typical seasonal stop log settings. Stop logs are typically set to lower water levels in December in preparation for the February to March prescribed burn, then returned to their original settings after the prescribed burn. The 2020 stop log and site hydrology data is contained in Appendix I.

Table 1 – Typical Seasonal Stop Log Settings

Wetland Cell	June – November	December – May
E-1	7:5	7
E-2	7:7	7:5
E-3	7:7	7:5
E-4	7:5	7
E-5	7:5	7
W-1	7:7	7
W-2	N/A	N/A

3.4 – Historical Herbicide Applications

ECO, Inc. routinely develops specifications for the types of herbicides and adjuvants that are applied for each targeted invasive plant species group. Use of broad-spectrum herbicides are avoided when practicable, but used on occasion. Herbicide applications are made following label recommendations and are not directly applied to standing surface water except when an EPA-approved Aquatic Site herbicide is used. Typically, graminicides are applied for control of invasive grass species and broadleaf-specific herbicides are applied for invasive broadleaf and woody plants. The majority of herbicide applications are made via backpack sprayers and/or UTV mounted spray equipment.

Historically, the predominate recurring and persistent non-native and invasive grasses to which herbicides have been applied, include:

- **tall fescue**
- **Johnsongrass**
- **small carpetgrass**
- **southern crabgrass**
- **smooth crabgrass**
- **Bermuda grass**

During pre-construction and early post-construction periods (pre-2006 through 2010) the very invasive tall fescue was the dominant plant species because the site was previously a hay pasture. Tall fescue has a wetland indicator status of FAC- and is capable of dominating wet meadow areas, significantly reducing native plant species richness. It is a cool season grass and actively begins photosynthesis very early in the growing season. It goes dormant during hot dry weather and actively grows in the fall even after several killing frosts. This provides an advantage in vegetation management since the fescue can have herbicide effectively applied at a time when native plant species are dormant. It has been observed that three to four weeks after the late winter/early spring burns are a critical time to apply herbicides on the fescue.

Since its original discovery in 2015 in the gas line easement west of wetland cell E-4, small carpetgrass has quickly become one of the most aggressive invasive species at the site and has now been observed as occurring in all of the wetland cells. Due to the nature of small carpet grass and its delayed growth cycle it is hard to observe new patches under the tall dense vegetation around it until it is already well established making it extremely hard to treat all of the individuals before going to seed in mid September. Historically, small carpetgrass has been treated with the herbicide Section 2EC (Clethodim 26.4%), which is a selective post-emergence herbicide used for the control of annual and perennial grasses. Monitoring events during 2020 showed a positive decrease in the densities of small carpet grass as compared to previous years. A positive sign that the herbicide concentration being used it working at controlling and effectively killing the species.

Recurring and persistent non-native and invasive broadleaf forbs to which herbicides have been applied, include:

- **sericea lespedeza**
- **white sweet clover**
- **curly dock**
- **bitter dock**
- **nodding thistle**
- **Japanese honeysuckle**
- **wooly mullein (*Verbascum Thapsus*)**
- **narrowleaf cattail**
- **Queen Anne's lace**
- **common thistle**
- **yellow rocket (*Barbarea vulgaris*)**

Herbicide applications to woody plant species have primarily focused upon black willow, green ash, honey locust (*Gleditsia triacanthos*), and persimmon (*Diospyros virginiana*). Complete eradication of these woody plants has been part of an ongoing vegetation management objective, but is a difficult task to achieve. In the past, selected stands of black willows have routinely been cut, with cut stumps sprayed with herbicide for the intent of a complete kill of individual trees; while green ash and persimmon saplings have had foliar and basal bark applications of herbicides.

Historically, the predominate recurring and persistent non-native and/or invasive woody plants to which herbicides have been applied, include:

- **black willow**
- **green ash**
- **persimmon**
- **callery pear**
- **honey locust**
- **winged elm (*Ulmus alata*)**
- **American elm (*Ulmus americana*)**
- **multiflora rose**
- **Chinese privet**
- **bush honeysuckle**
- **Himalayan blackberry**

During 2016 monitoring activities, it became more apparent to ECO, Inc. that the rate of black willow, persimmon, honey locust, and green ash stand expansion exceeded the rate of black willow cutting/spraying/eradication. As a part of developing the 2016 Woolsey Wet Prairie Adaptive Management Strategy & Monitoring Report No. 10, ECO, Inc. reviewed tree density information and recommended a more aggressive tree removal strategy to control woody plant succession. The increase in woody plant density was problematic, because the woody growth outcompetes the native prairie forbs and grasses, resulting in a reduction in both species diversity and density of the desirable native plants in all seven wetland cells. Prairie Ecosystems are level or slightly undulating, mostly treeless tracts of land, dominated by coarse grasses, forbs, and shrubs,

rather than trees, as the dominant vegetation type. The expanding stands of woody vegetation posed a threat to the prairie ecosystem character of the site to the point that if allowed to continue, the site would eventually become a forest.

The 2019 and 2020 growing seasons showed continued signs of success with the return of many of the rare ANHC tracking species that had previously been reduced by the woody vegetation, plus a continued increase in herbaceous plant species. The densities of the woody species outside of the “Designated Tree Preservation Areas” have been reduced to a much more manageable level of control. The practice of controlling woody vegetation outside of the Designated Tree Preservation Areas will be continued through the 2021 adaptive management year.

4.0 – RECOMMENDED ADAPTIVE MANAGEMENT ACTIVITIES FOR 2021

Based upon observations made during the 2020 and previous growing seasons, the anticipated general adaptive management activities for 2021 are contained in Table 2 below.

Table 2 – 2021 Woolsey Wet Prairie Adaptive Management Tentative Schedule*

General Timeframe	Activity
January through March	January Annual Adaptive Management Conference with management team and City of Fayetteville
	Final adjustments to fire line if necessary after Burn Contractor inspection
	Adjust stop logs to reduce water retention in all wetland cells in preparation for the prescribed burn, as needed.
	Conduct prescribed burn
	Reset stop logs to retain more water in wetland cells after prescribed burn.
	Observe re-emergence of tall fescue in 3-4 weeks after burn and apply Section 2-EC when tall fescue is 4-6 inches tall. Spray tall fescue with after prescribed burn and before native plants come out of dormancy.
March through April	Observe site for the emergence of yellow rocket, bull thistle, burdock, bush honeysuckle, and Japanese honeysuckle. Treat with PastureGard HL.
	Select wetland depressions for herbicide applications to native and non-native plants in order to restore mudflat microhabitats for shorebirds.
Late-March through September	Observe site for emergence of woody vegetation, including Himalayan blackberry, black willow saplings, green ash saplings, callery pear saplings, and honey locust saplings, all of which are to be treated using basal bark spray with Remedy Ultra for terrestrial and dry areas, or Renovate 3 for areas of standing water.
	Spray sericea lespedeza with PastureGard HL.
	Spray small carpetgrass with Section 2EC
May through September	Evaluate site for presence of yellow rocket; control via top-cutting/hand pulling/spray with Remedy Ultra.
	Observe site for emergence of white sweet clover (previously observed in Wetland Cells E-1, E-2, and E-3 along the inner portions of the berms); treat with PastureGard HL.
June through September	Adjacent (west and north) fescue fields to be mowed before tall fescue goes to seed
	Spot spray Johnsongrass with Outrider.
	Hand pull curly dock & Queen Anne’s lace.

* The appropriate concentrations and application rates of herbicides summarized in the Appendix II Woolsey Wet Prairie Sanctuary Herbicide Mixing/Application Rates should be followed.

5.0 – 2021 EXISTING SURPLUS WETLAND MITIGATION CREDITS

5.1 – Background and Overview

Subsequent to construction and initial adaptive management of Woolsey Wet Prairie, ECO, Inc. determined that 94.47 mitigation credits had been generated, producing a surplus of 20.90 credits above the required 73.57 wetland mitigation credits required by the COE Section 404 permit. ECO, Inc. and the City of Fayetteville met with the COE in mid-2013 to discuss the use of surplus wetland credits for city infrastructure projects that required wetland compensatory mitigation. On September 30, 2013, the City of Fayetteville received approval from the COE to use the 20.90 surplus wetland credits for impacts to wetlands caused by municipal projects within the Illinois River Watershed 8-digit Hydrologic Unit Code (HUC) watershed (11110103), but the City would not be allowed to sell the surplus credits.

Consequently, the City of Fayetteville's surplus wetland credits in essence serve as a mitigation bank where improvements to wetland ecological function and value provide an ecological gain and are available to be used to meet compensatory mitigation requirements for city projects that permanently alter wetlands. More specifically, these surplus credits serve as what is known as a Single-Client Mitigation Bank, or a bank for which the sponsor is also the principal credit user or client.

To date, in 2014, the COE authorized the use of a portion of the Woolsey Wet Prairie surplus wetland credits to offset 0.31 acres of permanent alterations to wetlands from the construction of an extension to Van Ashe Drive (COE Project No. 2012-00525). Consequently, the City of Fayetteville Transportation Division Van Ashe Drive project deducted 2.94 credits from the Woolsey Wet Prairie 20.90 surplus wetland credits, leaving a balance of 17.96 surplus credits.

In 2015, the COE authorized the use of a portion of the Woolsey Wet Prairie surplus wetland credits to offset 0.192 acres of permanent alterations to wetlands from the construction of the Clabber Creek Recreational Trail. Consequently, the City of Fayetteville Clabber Creek Recreational Trail project deducted 3.14 credits from the remaining 17.96 Woolsey Wet Prairie surplus wetland credits, leaving a balance of 14.82 surplus credits.

The City of Fayetteville did not use any surplus wetland credits in 2016, 2017, 2018, or 2019.

In 2020, the COE authorized the use of a portion of the Woolsey Wet Prairie surplus wetland credits to offset 0.153 acres of permanent alterations to wetlands from the construction of another Clabber Creek Recreational Trail. Consequently, the City of Fayetteville Clabber Creek Recreational Trail project deducted 2.50 credits from the remaining 14.82 Woolsey Wet Prairie surplus wetland credits, leaving a balance of 12.32 surplus credits. Although the surplus wetland credit balance is not a large number, having them available for small municipal infrastructure improvements projects that impact wetlands has saved the City of Fayetteville in excess of tens of thousands of dollars.

6.0 – Academic Research and Guidance

In 2017 and 2018, there was an increased interest by the academic community in conducting scientific research at Woolsey Wet Prairie. ECO, Inc. has met with various professors and students to provide guidance to them concerning minimizing adverse impacts to the site during their fieldwork. To date, low impact studies have been conducted, including insect trapping, bird surveys, and herpetological surveys that have produced some amazing results.

In order to further aid in educating academia interested in conducting research at the site, ECO, Inc. developed a document entitled "***Guidance For Conducting Scientific Research At Woolsey Wet Prairie Sanctuary***" that was been approved by the COE and City of Fayetteville. This document can be found on the ECO, Inc. Woolsey Wet Prairie Facebook page or the ECO, Inc. website. As a part of the guidance document, ECO, Inc. developed a research project application form for prospective academic projects to complete and subsequently be approved by the City of Fayetteville and ECO, Inc.

ECO, Inc. and the City of Fayetteville welcomes academic research projects at the site and has tried to make the approval process as streamlined as possible. It became necessary to develop this guidance to inform those who desire to conduct scientific research that Woolsey Wet Prairie Sanctuary is like no other City-owned property. It is a wetland compensatory mitigation site that is deed restricted in perpetuity with specific restrictions and is authorized under a Section 404 permit issued by the COE to the City of Fayetteville that will never expire. The City is required to comply with the terms and conditions of the COE permit and deed restrictions for protecting the site.

8.0 – Appendices

Appendix I
2020 Stop Log Data For
Hydrological Controls In Wetland Cells

Year		2020 Woolsey Wet-Prairie Annual Surface Water Monitoring Table																																													
Month				Jan		Feb		Mar		Apr		May		Jun		Jul		Aug		Sep		Oct		Nov		Dec		ANN																			
Precip.	ACT	NRM	DIFF	6.4	2.8	3.6	3.4	2.8	0.6	8.0	4.0	4.0	5.3	4.6	0.7	9.0	6.0	3.0	1.2	5.0	-3.8	5.3	3.5	1.8	1.9	3.2	-1.3	2.7	4.8	-2.1	7.0	4.3	2.7	2.4	4.2	-1.8	1.0	3.2	-2.3	53.6	48.5	5.1					
Wetland Cells	E5	Surface H ₂ O	x				x			x			x			x			x		-		-		-		x		x		x								9	of	12						
		WOF Ret.	X				X			X			X			X			X		-		-		-		-		-		-								5	of	12						
	12" Max	Log Combo	0:				0:			7:			7:			5:5			0:		0:		0:		0:		0:		0:		0:																
		Height (in.)	0				0			7			7			10			0		0		0		0		0		0		0		5														
	E4	Pond Ret.	x				x			x			x			x			x		x		x		x		x		x		x										12	of	12				
		S. H ₂ O	x				x			x			x			x			x		-		-		-		x		x		x										9	of	12				
	19" Max	WOF Ret.	X				X			X			X			X			-		-		-		-		-		-		-												5	of	12		
		L.C.	0:				0:			7:			7:			7:			7:		7:		7:		7:		7:		7:		5:																
		H. (in.)	0				0			7			7			7			7		7		7		7		7		7		7		5														
		E3	S. H ₂ O	x				x			x			x			x			x		-		-		-		-		-		-										7	of	12			
	21" Max	WOF Ret.	X				X			X			X			X			-		-		-		-		-		-		-												5	of	12		
		L.C.	0:				0:			7:			7:			5:5			5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5						
		H. (in.)	0				0			7			7			10			10		10		10		10		10		10		10		10		5												
		E1	S. H ₂ O	x				x			x			x			x			x		-		-		-		-		-		-											7	of	12		
	21" Max	WOF Ret.	X				X			X			X			X			-		-		-		-		-		-		-													5	of	12	
		L.C.	0:				0:			7:			7:			5:5			5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5						
		H. (in.)	0				0			7			7			10			10		10		10		10		10		10		10		10		5												
		E2	S. H ₂ O	x				x			x			x			x			x		-		-		-		-		-		-											7	of	12		
	21" Max	WOF Ret.	X				X			X			X			X			X		-		-		-		-		-		-													6	of	12	
		L.C.	0:				0:			7:			7:			5:5			5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5						
	H. (in.)	0				0			7			7			10			10		10		10		10		10		10		10		10		5													
W1	S. H ₂ O	x				x			x			x			x			x		x		-		-		-		x		x												10	of	12			
	WOF Ret.	X				X			X			X			X			-		-		-		-		-		-		-												5	of	12			
17" Max	L.C.	0:				0:			7:			7:			5:5			5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5		5:5							
	H. (in.)	0				0			7			7			10			10		10		10		10		10		10		10		10		5													
W2	Pond Ret.	x				x			x			x			x			x		x		x		x		x		x		x													12	of	12		
	S. H ₂ O	x				x			x			x			x			x		-		-		-		-		x		x												9	of	12			
	Max Ret.	X				X			X			X			X			X		-		-		-		-		-		-													6	of	12		
	L.C.	NA				NA			NA			NA			NA			NA		NA		NA		NA		NA		NA		NA		NA		NA		NA		NA		NA							
	H. (in.)	NA				NA			NA			NA			NA			NA		NA		NA		NA		NA		NA		NA		NA		NA		NA		NA		NA							


Appendix II
Woolsey Wet Prairie Sanctuary
Herbicide Mixing/Application Rates

WOOLSEY WET PRAIRIE SANCTUARY HERBICIDE MIXING/APPLICATION RATES

Herbicide	Target Species	Label Mixing Rate for Spot Spray <u>Control</u> of Target Plant Species
Remedy Ultra (60.45% triclopyr)	Broadleaf Weeds	3.2 fl. oz. Remedy Ultra + 0.68 fl. oz. surfactant + 124.12 fl. oz. water = 128 fl. oz. = 1 gal. of 2.5% mixture
Remedy Ultra	Woody Plants Cut stump/basal bark	32 fl. oz. Remedy Ultra + 96 fl. oz. Oil Carrier (in lieu of water + surfactant) = 1 gal. of 25% mixture Can be mixed with a penetrant such as Cide-kick in enough oil to make the desired volume of spray mixture to penetrate the cuticle of the woody plant, allowing more herbicide to translocate to the root system. Oil carrier should be a petroleum-based oil or a vegetable oil concentrate. DO NOT use diesel or kerosene at Woolsey Wet Prairie. For basil bark spraying stems < 6 inches in diameter, thoroughly wet the base and root collar of all stems to a height of 12 to 15 inches, but not to the point of runoff.
Remedy Ultra	Woody Plants Foliar	6.8 fl. oz. Remedy Ultra + 0.68 fl. oz. surfactant + 128 fl. oz. water = 1.06 gal. of 5.0% mixture Depending upon the size and density of the woody plants, apply sufficient spray volume to thoroughly wet all leaves, stems, and root collars. This concentration works best for regrowth less than 4 feet tall.
PastureGard HL (45% triclopyr + 15.56% fluroxypyr)	Broadleaf Weeds; Herbicide of choice for sericea lespedeza and white sweet clover	1.35 fl. oz. PastureGard HL + 0.68 fl. oz. surfactant + 125.97 fl. oz. water = 128 fl. oz. = 1 gal. = 1.0% mixture
PastureGard HL	Woody Plants Cut stump/basal bark	32 fl. oz. PastureGard HL + 96 fl. oz. Oil Carrier (in lieu of water + surfactant) = 1 gal. of 25% mixture Can be mixed with a penetrant such as Cide-kick in enough oil to make the desired volume of spray mixture to penetrate the cuticle of the woody plant, allowing more herbicide to translocate to the root system. Oil carrier should be a petroleum-based oil or a vegetable oil concentrate. DO NOT use diesel or kerosene at Woolsey Wet Prairie. For basil bark spraying stems < 6 inches in diameter, thoroughly wet the base and root collar of all stems to a height of 12 to 15 inches, but not to the point of runoff.
Renovate® 3 (44.4% trichlopyr)	Woody Plants: Allowable to spray over water basal bark or cut surface treatment	Can be applied either undiluted or diluted in a 1 to 1 ratio with water and sprayed where the mixture will completely surround the tree trunk. Cide-Kick to be mixed with Renovate 3 as a penetrant to increase bark penetration.
Roundup Pro Concentrate (50.2% glyphosate)	Broad-spectrum non-selective chemical used to control both grasses and forbs.	Use a 1.6% solution for harder to control perennials such as Bermuda grass, dock, thistle, sericea, etc. This would be the equivalent of 2.1 fl. oz. Roundup Pro Concentrate in 125.9 fl. oz. of water to make one gallon of spray mix. Roundup Pro Concentrate already contains a 13% surfactant, so none should be added.
Section 2EC (26.4% clethodim)	Grasses; best herbicide for tall fescue, especially in moist soils because it will not harm rushes.	Mix 1.2 fl. oz. Section 2 EC + 1.66 fl. oz. Superb HC surfactant + 0.4 fl. oz. Mystic HC dye + 124.74 fl. oz. water = 1 gal. of 1% mixture
Outrider (75% Sulfosulfuron)	Grasses; best herbicide for Johnsongrass in upland habitat.	Mix 0.25 fl. oz. Outrider + 0.1 fl. oz. nonionic surfactant + 127.65 fl. oz. water = 1 gal. of 0.2% mixture

Appendix III
Tracking Species
Aerial Images

Wetland Cell E-1 2020 Tracking Species Aerial Image



LEGEND:


O – OPAQUE PRAIRIE SEDGE (*CAREX OPACA*)

AG – CLUSTER SEDGE (*CAREX AGGREGATA*)

AR – ARKANSAS SEDGE (*CAREX ARKANSANA*)



Wetland Cell E-2 2020 Tracking Species Aerial Image



LEGEND:

O – OPAQUE PRAIRIE SEDGE (*CAREX OPACA*)

AR – ARKANSAS SEDGE (*CAREX ARKANSANA*)

AG – CLUSTER SEDGE (*CAREX AGGREGATA*)

W – WOLF’S SPIKERUSH (*ELEOCHARIS WOLFII*)

A DASHED CIRCLE INDICATES A LARGE PATCH OF THE TRACKING SPECIES INDICATED IN THE ASSOCIATED TEXT BOX.



Wetland Cell E-3 2020 Tracking Species Aerial Image



LEGEND:

O – OPAQUE PRAIRIE SEDGE (*CAREX OPACA*)

AR – ARKANSAS SEDGE (*CAREX ARKANSANA*)


AG – CLUSTER SEDGE (*CAREX AGGREGATA*)

W – WOLF’S SPIKERUSH (*ELEOCHARIS WOLFII*)

A DASHED CIRCLE INDICATES A LARGE PATCH OF THE TRACKING SPECIES INDICATED IN THE ASSOCIATED TEXT BOX.



Wetland Cell E-4 2020 Tracking Species Aerial Image




LEGEND:

- O – OPAQUE PRAIRIE SEDGE (*CAREX OPACA*)
- AR – ARKANSAS SEDGE (*CAREX ARKANSANA*)
- AG – CLUSTER SEDGE (*CAREX AGGREGATA*)
- W – WOLF’S SPIKERUSH (*ELEOCHARIS WOLFII*)
- CP – WOOLLY SEDGE (*CAREX PELLITA*)
- CR – REVERCHON’S HAWTHORN (*CRATEGUS REVERCHONII*)

A DASHED CIRCLE INDICATES A LARGE PATCH OF THE TRACKING SPECIES INDICATED IN THE ASSOCIATED TEXT BOX.



Wetland Cell E-5 2020 Tracking Species Aerial Image



LEGEND:

- O – OPAQUE PRAIRIE SEDGE (*CAREX OPACA*)
- AR – ARKANSAS SEDGE (*CAREX ARKANSANA*)
- AG – CLUSTER SEDGE (*CAREX AGGREGATA*)
- W – WOLF’S SPIKERUSH (*ELEOCHARIS WOLFII*)
- CS – POINTED SEDGE (*CAREX SCOPARIA*)
- CP – WOOLLY SEDGE (*CAREX PELLITA*)
- R – TALL-HORNED BEAKSEEDGE (*RHYNCOSEPORA MACROSTACHYA*)

A DASHED CIRCLE INDICATES A LARGE PATCH OF THE TRACKING SPECIES INDICATED IN THE ASSOCIATED TEXT BOX.



Gas Line Easement 2020 Tracking Species Aerial Image



Wetland Cell W-1 2020 Tracking Species Aerial Image




LEGEND:

- O – OPAQUE PRAIRIE SEDGE (*CAREX OPACA*)
- AR – ARKANSAS SEDGE (*CAREX ARKANSANA*)
- AG – CLUSTER SEDGE (*CAREX AGGREGATA*)
- W – WOLF’S SPIKERUSH (*ELEOCHARIS WOLFII*)
- CS – POINTED SEDGE (*CAREX SCOPARIA*)
- R – TALL-HORNED BEAKSEGE (*RHYNOSPORA MACROSTACHYA*)
- F – HAMMOCK SEDGE (*CAREX FISSA*)
- M – MEXICAN WHITE-SAGE (*ARTEMISIA LUDOVICIANA VAR MEXICANA*)



Wetland Cell W-2 2020 Tracking Species Aerial Image




LEGEND:

- O – OPAQUE PRAIRIE SEDGE (*CAREX OPACA*)
 - AR – ARKANSAS SEDGE (*CAREX ARKANSANA*)
 - AG – CLUSTER SEDGE (*CAREX AGGREGATA*)
 - W – WOLF’S SPIKERUSH (*ELEOCHARIS WOLFII*)
 - CS – POINTED SEDGE (*CAREX SCOPARIA*)
 - CP – WOOLLY SEDGE (*CAREX PELLITA*)
 - R – TALL-HORNED BEAKSEGE (*RHYNOSPORA MACROSTACHYA*)
 - F – HAMMOCK SEDGE (*CAREX FISSA*)
 - CR – REVERCHON’S HAWTHORN (*CRATEGUS REVERCHONII*)
 - M – MEXICAN WHITE-SAGE (*ARTEMISIA LUDOVICIANA VAR MEXICANA*)
- A DASHED CIRCLE INDICATES A LARGE PATCH OF THE TRACKING SPECIES INDICATED IN THE ASSOCIATED TEXT BOX.



Appendix IV
Invasive Species
Aerial Images

2020 Invasive Species Monitoring Map Wetland Cell E-1



LEGEND:

RED – JUNE 2020
GROWING SEASON MONITORING EVENT

BLUE – AUGUST 2020
POST GROWING SEASON MONITORING EVENT

L – SERICEA LESPEDEZA (*LESPEDEZA CUNEATA*)

M – WHITE SWEET CLOVER (*MELILOTUS ALBUS*)

A – SMALL CARPETGRASS (*ARTHRAOXON HISPIDUS*)

C – CALLERY PEAR (*PYRUS CALLERYANA*)

A DASHED CIRCLE INDICATES A LARGE
PATCH OF THE INVASIVE SPECIES INDICATED
IN THE ASSOCIATED TEXT BOX.

A SOLID CIRCLE AROUND A LETTER
REPRESENTS THAT THE INVASIVE SPECIES
WAS OBSERVED AT THE SAME LOCATION AS
THE PREVIOUS MONITORING EVENT.



2020 Invasive Species Monitoring Map Wetland Cell E-2



LEGEND:

RED – JUNE 2020
GROWING SEASON MONITORING EVENT

BLUE – AUGUST 2020
POST GROWING SEASON MONITORING EVENT

L – SERICEA LESPEDEZA (*LESPEDEZA CUNEATA*)

M – WHITE SWEET CLOVER (*MELILOTUS ALBUS*)

A – SMALL CARPETGRASS (*ARTHRAOXON HISPIDUS*)

C – CALLERY PEAR (*PYRUS CALLERYANA*)

R – MULTIFLORA ROSE (*ROSA MULTIFLORA*)

J – JAPANESE HONEYSUCKLE (*LONICERA JAPONICA*)

T – NARROW-LEAF CATTAIL (*TYPHA ANGUSTIFOLIA*)

A DASHED CIRCLE INDICATES A LARGE
PATCH OF THE INVASIVE SPECIES INDICATED
IN THE ASSOCIATED TEXT BOX.

A SOLID CIRCLE AROUND A LETTER
REPRESENTS THAT THE INVASIVE SPECIES
WAS OBSERVED AT THE SAME LOCATION AS
THE PREVIOUS MONITORING EVENT.



2020 Invasive Species Monitoring Map Wetland Cell E-3




LEGEND:

- RED – JUNE 2020
GROWING SEASON MONITORING EVENT**
- BLUE – AUGUST 2020
POST GROWING SEASON MONITORING EVENT**
- L – SERICEA LESPEDEZA (*LESPEDEZA CUNEATA*)**
- M – WHITE SWEET CLOVER (*MELILOTUS ALBUS*)**
- A – SMALL CARPETGRASS (*ARTHRAOXON HISPIDUS*)**
- C – CALLERY PEAR (*PYRUS CALLERYANA*)**
- A DASHED CIRCLE INDICATES A LARGE
PATCH OF THE INVASIVE SPECIES INDICATED
IN THE ASSOCIATED TEXT BOX.**
- A SOLID CIRCLE AROUND A LETTER
REPRESENTS THAT THE INVASIVE SPECIES
WAS OBSERVED AT THE SAME LOCATION AS
THE PREVIOUS MONITORING EVENT.**



2020 Invasive Species Monitoring Map Wetland Cell E-4



LEGEND:

RED – JUNE 2020
GROWING SEASON MONITORING EVENT

BLUE – AUGUST 2020
POST GROWING SEASON MONITORING EVENT

L – SERICEA LESPEDEZA (*LESPEDEZA CUNEATA*)

M – WHITE SWEET CLOVER (*MELILOTUS ALBUS*)

A – SMALL CARPETGRASS (*ARTHRAOXON HISPIDUS*)

R – MULTIFLORA ROSE (*ROSA MULTIFLORA*)

H – HIMALAYAN BLACKBERRY (*RUBUS SERISSIMUS*)


V – JAPANESE STILTGRASS (*MICROSTEGIUM
VIMINEUM*)

A DASHED CIRCLE INDICATES A LARGE
PATCH OF THE INVASIVE SPECIES INDICATED
IN THE ASSOCIATED TEXT BOX.

A SOLID CIRCLE AROUND A LETTER
REPRESENTS THAT THE INVASIVE SPECIES
WAS OBSERVED AT THE SAME LOCATION AS
THE PREVIOUS MONITORING EVENT.



2020 Invasive Species Monitoring Map Wetland Cell E-5



LEGEND:

RED – JUNE 2020
GROWING SEASON MONITORING EVENT

BLUE – AUGUST 2020
POST GROWING SEASON MONITORING EVENT

L – SERICEA LESPEDEZA (*LESPEDEZA CUNEATA*)

A – SMALL CARPETGRASS (*ARTHRAON HISPIDUS*)

J – JAPANESE HONEYSUCKLE (*LONICERA JAPONICA*)


T – NARROW-LEAF CATTAIL (*TYPHA ANGUSTIFOLIA*)

A DASHED CIRCLE INDICATES A LARGE
PATCH OF THE INVASIVE SPECIES INDICATED
IN THE ASSOCIATED TEXT BOX.


A SOLID CIRCLE AROUND A LETTER
REPRESENTS THAT THE INVASIVE SPECIES
WAS OBSERVED AT THE SAME LOCATION AS
THE PREVIOUS MONITORING EVENT.



2020 Invasive Species Monitoring Map Gas Line Easement



2020 Invasive Species Monitoring Map Wetland Cell W-1




LEGEND:

- RED – JUNE 2020
GROWING SEASON MONITORING EVENT
- BLUE – AUGUST 2020
POST GROWING SEASON MONITORING EVENT
- L – SERICEA LESPEDEZA (*LESPEDEZA CUNEATA*)
- M – WHITE SWEET CLOVER (*MELILOTUS ALBUS*)
- A – SMALL CARPETGRASS (*ARTHRAOXON HISPIDUS*)
- C – CALLERY PEAR (*PYRUS CALLERYANA*)
- H – HIMALAYAN BLACKBERRY (*RUBUS SERISSIMUS*)
- X – HYBRID RUBUS SPP.
- J – JAPANESE HONEYSUCKLE (*LONICERA JAPONICA*)
- T – NARROW-LEAF CATTAIL (*TYPHA ANGUSTIFOLIA*)
- A DASHED CIRCLE INDICATES A LARGE
PATCH OF THE INVASIVE SPECIES INDICATED
IN THE ASSOCIATED TEXT BOX.
- A SOLID CIRCLE AROUND A LETTER
REPRESENTS THAT THE INVASIVE SPECIES
WAS OBSERVED AT THE SAME LOCATION AS
THE PREVIOUS MONITORING EVENT.



2020 Invasive Species Monitoring Map Wetland Cell W-2



LEGEND:

RED – JUNE 2020
GROWING SEASON MONITORING EVENT

BLUE – AUGUST 2020
POST GROWING SEASON MONITORING EVENT

- L – SERICEA LESPEDEZA (*LESPEDEZA CUNEATA*)
- A – SMALL CARPETGRASS (*ARTHRAON HISPIDUS*)
- C – CALLERY PEAR (*PYRUS CALLERYANA*)
- H – HIMALAYAN BLACKBERRY (*RUBUS SERISSIMUS*)
- X – HYBRID RUBUS SPP.
- J – JAPANESE HONEYSUCKLE (*LONICERA JAPONICA*)
- T – NARROW-LEAF CATTAIL (*TYPHA ANGUSTIFOLIA*)
- S – JOHNSON GRASS (*SORGHUM HALEPENSE*)

A DASHED CIRCLE INDICATES A LARGE
PATCH OF THE INVASIVE SPECIES INDICATED
IN THE ASSOCIATED TEXT BOX.

A SOLID CIRCLE AROUND A LETTER
REPRESENTS THAT THE INVASIVE SPECIES
WAS OBSERVED AT THE SAME LOCATION AS
THE PREVIOUS MONITORING EVENT.



Appendix V
Woolsey Wet Prairie
2020 Master Plant Species List

MASTER PLANT SPECIES LIST FOR WOOLSEY WET PRAIRIE - updated December 2020 - 496 taxa

SCIENTIFIC NAME	WETLAND INDICATOR STATUS	CODE	COMMON NAME	FAMILY	STRATA	SOURCE CODE
<i>Abutilon theophrastii</i> *	FACU-	ABUT THEO	pie-maker	MALVACEAE	herb	5
<i>Acalypha gracilens</i>	no data	ACAL GRAC	copperleaf	EUPHORBIACEAE	herb	5
<i>Acalypha virginica</i>	FACU-	ACAL VIRG	Virginia copperleaf	EUPHORBIACEAE	herb	3
<i>Acer negundo</i>	FACW	ACER NEGU	boxelder	ACERACEAE	tree/sapling	5
<i>Acer saccharinum</i>	FACW	ACER SACC	silver maple	ACERACEAE	tree/sapling	9
<i>Achillea millefolium</i>	FACU	ACHI MILL	yarrow	ASTERACEAE	herb	12
<i>Agalinis fasciculata</i>	FAC	AGAL FASC	gerardia	SCROPHULARIACEAE	herb	6
<i>Agrimonia parviflora</i> +	FACW	AGRI PARV	swamp agrimony	ROSACEAE	herb	21
<i>Agrostis gigantea</i> *	FACW	AGRO GIGA	redtop	POACEAE	herb	3
<i>Agrostis hyemalis</i>	FAC	AGRO HYEM	ticklegass	POACEAE	herb	5
<i>Ailanthus altissima</i> **	NI	AILA ALTI	tree-of-heaven	SIMAROUACEAE	tree/sapling	10
<i>Allium canadense</i> var. <i>canadense</i>	FACU	ALLI CANA CANA	wild onion	ALLIACEAE	herb	19
<i>Allium vineale</i> *	FACU-	ALLI VINE	field garlic	ALLIACEAE	herb	3
<i>Amaranthus</i> cf. <i>viridis</i> *	NO	AMAR VIRI	pigweed	AMARANTHACEAE	herb	4
<i>Amaranthus spinosus</i>	FACU	AMAR SPIN	spiny pigweed	AMARANTHACEAE	herb	3
<i>Ambrosia artemisiifolia</i>	FACU	AMBR ARTE	common ragweed	ASTERACEAE	herb	3
<i>Ambrosia bidentata</i>	no data	AMBR BIDE	lanceleaf ragweed	ASTERACEAE	herb	3
<i>Ambrosia trifida</i>	FAC	AMBR TRIF	giant ragweed	ASTERACEAE	herb	3
<i>Ammannia X coccinea</i>	FACW+	AMMA COCC	toothcup	LYTHRACEAE	herb	3
<i>Amorpha fruticosa</i> +	FACW	AMOR FRUT	false indigo bush	FABACEAE	shrub	21
<i>Ampelopsis cordata</i>	FAC+	AMPE CORD	heartleaf ampelopsis	VITACEAE	woody vine	10
<i>Anagallis minima</i>	FACW	ANAG MINI	chaffweed	PRIMULACEAE	herb	23
<i>Andropogon gerardii</i>	FAC	ANDR GERA	big bluestem	POACEAE	herb	3
<i>Andropogon glomeratus</i>	FACW+	ANDR GLOM	bushy bluestem	POACEAE	herb	11
<i>Andropogon hirsutior</i>	FACW	ANDR HIRS	hirsute bushy bluestem	POACEAE	herb	24
<i>Andropogon ternarius</i> Michx.	FACU	ANDR TERN	splitbeard bluestem	POACEAE	herb	32
<i>Andropogon virginicus</i>	FAC-	ANDR VIRG	broomsedge bluestem	POACEAE	herb	3
<i>Apios americana</i>	FACW	APIO AMER	groundnut	FABACEAE	herb	3
<i>Apocynum cannabinum</i>	FAC-	APOC CANN	Indian hemp	APOCYNACEAE	herb	3
<i>Arctium minus</i> *	FACU	ARCT MINU	burdock	ASTERACEAE	herb	19
<i>Arenaria serpyllifolia</i> var. <i>tenuior</i> *	FAC	AREN SERP TENU	thyme-leaved sandwort	CARYOPHYLLACEAE	herb	19
<i>Aristida dichotoma</i>	FACU	ARIS DICH	churchmouse three-awn	POACEAE	herb	8
<i>Aristida longespica</i> var. <i>longespica</i>	UPL	ARIS LONG	slimspike three-awn	POACEAE	herb	22
<i>Aristida oligantha</i>	no data	ARIS OLIG	three-awn	POACEAE	herb	8
<i>Arthraxon hispidus</i> **	FAC	ARTH HISP	small carpetgrass	POACEAE	herb	23
<i>Artemisia ludoviciana</i> var. <i>mexicana</i>	UPL	ARTE LUDO MEXI	Mexican white sage	ASTERACEAE	herb	31
<i>Asclepias amplexicaulis</i>	no data	ASCL AMPL	curly milkweed	ASCLEPIADACEAE	herb	3
<i>Asclepias longifolia</i> ssp. <i>hirtella</i>	UPL	ASCL HIRT	longleaf milkweed	ASCLEPIADACEAE	herb	3
<i>Asclepias incarnata</i>	OBL	ASCL INCA	swamp milkweed	ASCLEPIADACEAE	herb	30
<i>Asclepias viridis</i>	no data	ASCL VIRI	spider milkweed	ASCLEPIADACEAE	herb	3
<i>Baptisia alba</i> var. <i>macrophylla</i>	no data	BAPT ALBA	white false indigo	FABACEAE	herb	3
<i>Baptisia bracteata</i> var. <i>leucophaea</i>	no data	BAPT BRAC	cream false indigo	FABACEAE	herb	1
<i>Barbarea vulgaris</i> **	FAC	BARB VULG	yellow rocket	BRASSICACEAE	herb	5
<i>Bidens aristosa</i>	FACW	BIDE ARIS	tickseed sunflower	ASTERACEAE	herb	8
<i>Bidens frondosa</i>	FACW	BIDE FRON	tickseed sunflower	ASTERACEAE	herb	13

<i>Boehmeria cylindrica</i>	FACW	BOEH CYLI	false nettle	URTICACEAE	herb	27
<i>Boltonia asteroides</i>	FACW	BOLT ASTE	false aster	ASTERACEAE	herb	3
<i>Boltonia diffusa</i>	FAC	BOLT DIFF	doll's daisy	ASTERACEAE	herb	8
<i>Bradburia pilosa</i>	FACW	BRAD PILO	soft goldaster	ASTERACEAE	herb	32
<i>Brassica rapa</i> *	no data	BRAS RAPA	turnip	BRASSICACEAE	herb	5
<i>Bromus catharticus</i> *	no data	BROM CATH	rescue grass	POACEAE	herb	5
<i>Bromus hordeaceus</i> *	no data	BROM HORD	soft chess	POACEAE	herb	5
<i>Bromus inermis</i> *	no data	BROM INER	smooth broome	POACEAE	herb	5
<i>Bromus racemosus</i> **	no data	BROM RACE	bald brome	POACEAE	herb	5
<i>Bromus sterilis</i> **	no data	BROM STER	poverty brome	POACEAE	herb	19
<i>Bromus tectorum</i> *	no data	BROM TECT	cheatgrass	POACEAE	herb	5
<i>Callitriche heterophylla</i>	OBL	CALL HETE	water starwort	CALLITRICHACEAE	herb	5
<i>Callitriche terrestris</i>	FACW	CALL TERR	terrestrial water starwort	CALLITRICHACEAE	herb	19
<i>Campsis radicans</i>	FAC	CAMP RADI	trumpet creeper	BIGNONIACEAE	herb	10
<i>Capsella bursa-pastoris</i> *	FACU+	CAPS BURS	shepherd's purse	BRASSICACEAE	herb	5
<i>Cardamine parviflora</i> var. <i>arenicola</i>	FACU	CARD PARV AREN	small-flowered bittercress	BRASSICACEAE	herb	5
<i>Carduus nutans</i> **	no data	CARD NUTA	nodding thistle	ASTERACEAE	herb	4
<i>Carex aggregata</i>	no data	CARX AGGR	cluster sedge	CYPERACEAE	herb	19
<i>Carex amphibola</i>	FAC	CARX AMPH	a sedge	CYPERACEAE	herb	19
<i>Carex annectens</i>	FACW	CARX ANNE	a sedge	CYPERACEAE	herb	5
<i>Carex arkansana</i>	no data	CARX ARKA	Arkansas sedge	CYPERACEAE	herb	5
<i>Carex aureolensis</i>	no data	CARX AURE	a sedge	CYPERACEAE	herb	19
<i>Carex austrina</i>	no data	CARX AUST	a sedge	CYPERACEAE	herb	5
<i>Carex blanda</i>	FAC	CARX BLAN	a sedge	CYPERACEAE	herb	19
<i>Carex brevior</i>	OBL	CARX BREV	a sedge	CYPERACEAE	herb	5
<i>Carex bushii</i>	FACW	CARX BUSH	Bush's sedge	CYPERACEAE	herb	5
<i>Carex complanata</i>	FAC+	CARX COMP	a sedge	CYPERACEAE	herb	5
<i>Carex festucacea</i>	FACW	CARX FEST	a sedge	CYPERACEAE	herb	5
<i>Carex fissa</i>	FACW+	CARX FISS	hammock sedge	CYPERACEAE	herb	5
<i>Carex flaccosperma</i>	FAC+	CARX FLAC	a sedge	CYPERACEAE	herb	9
<i>Carex frankii</i>	OBL	CARX FRAN	Frank's sedge	CYPERACEAE	herb	5
<i>Carex glaucoidea</i>	no data	CARX GLAU	blue sedge	CYPERACEAE	herb	15
<i>Carex granularis</i>	FACW	CARX GRAN	granular sedge	CYPERACEAE	herb	5
<i>Carex grisea</i>	FACU	CARX GRIS	inflated narrow-leaf sedge	CYPERACEAE	herb	26
<i>Carex hirsutella</i>	no data	CARX HIRS	a sedge	CYPERACEAE	herb	5
<i>Carex leavenworthii</i>	no data	CARX LEAV	Leavenworth's sedge	CYPERACEAE	herb	5
<i>Carex meadii</i>	FAC	CARX MEAD	Mead's sedge	CYPERACEAE	herb	7
<i>Carex molestiformis</i>	no data	CARX MOLE	frightful sedge	CYPERACEAE	herb	26
<i>Carex oklahomensis</i>	OBL	CARX OKLA	Oklahoma sedge	CYPERACEAE	herb	3
<i>Carex opaca</i>	no data	CARX OPAC	opaque prairie sedge	CYPERACEAE	herb	5
<i>Carex pellita</i>	OBL	CARX PELL	woolly sedge	CYPERACEAE	herb	5
<i>Carex retroflexa</i>	no data	CARX RETR	a sedge	CYPERACEAE	herb	5
<i>Carex scoparia</i>	FACW	CARX SCOP	pointed sedge	CYPERACEAE	herb	16
<i>Carex shortiana</i>	FACW	CARX SHOR	Short's sedge	CYPERACEAE	herb	14
<i>Carex triangularis</i>	FACW	CARX TRIA	eastern fox sedge	CYPERACEAE	herb	31
<i>Carex vulpinoidea</i>	OBL	CARX VULP	foxtail sedge	CYPERACEAE	herb	3
<i>Carya illinoensis</i>	FACU	CARY ILLI	pecan	JUGLANDACEAE	tree/sapling	20
<i>Catalpa bignonioides</i>	FAC-	CATA BIGN	catalpa	BIGNONIACEAE	tree/sapling	3
<i>Celtis laevigata</i>	FACW	CELT LAEV	sugarberry	CELTIDACEAE	tree/sapling	20
<i>Celtis occidentalis</i>	FACU	CELT OCCI	hackberry	CELTIDACEAE	tree/sapling	8
<i>Cephalanthus occidentalis</i>	OBL	CEPH OCCI	buttonbush	RUBIACEAE	shrub	3
<i>Cerastium fontanum</i> ssp. <i>vulgare</i> *	no data	CERA FONT VULG	chickweed	CARYOPHYLLACEAE	herb	19

<i>Cerastium pumilum</i> *	no data	CERA PUMI	chickweed	CARYOPHYLLACEAE	herb	4
<i>Ceratophyllum demersum</i>	OBL	CERA DEME	coontail	CERATOPHYLLACEAE	herb	10
<i>Chamaesyce humistrata</i>	FAC	CHAM HUMI	spreading sandmat	EUPHORBIACEAE	herb	27
<i>Chamaesyce maculata</i>	no data	CHAM MACU	spotted spurge	EUPHORBIACEAE	herb	3
<i>Chamaesyce nutans</i>	FACU	CHAM NUTA	spurge	EUPHORBIACEAE	herb	8
<i>Chasmanthium latifolium</i>	FACU	CHAS LATI	river oats	POACEAE	herb	23
<i>Chenopodium album</i>	FAC-	CHEN ALBU	lamb's quarters	CHENOPODIACEAE	herb	3
<i>Cicuta maculata</i>	OBL	CICU MACU	water hemlock	APIACEAE	herb	5
<i>Cirsium altissimum</i>	no data	CIRS ALTI	tall thistle	ASTERACEAE	herb	19
<i>Cirsium vulgare</i> **	FAC	CIRS VULG	common thistle	ASTERACEAE	herb	8
<i>Claytonia virginica</i>	FAC	CLAY VIRG	spring beauty	PORTULACACEAE	herb	18
<i>Cocculus carolinus</i>	FAC	COCC CARO	Carolina snailseed	MENISPERMACEAE	herb	20
<i>Conium maculatum</i> *	FACW	CONI MACU	poison hemlock	APIACEAE	herb	7
<i>Conoclinium coelestinum</i>	FAC	CONO COEL	blue mistflower	ASTERACEAE	herb	31
<i>Conyza canadensis</i>	FACU	CONY CANA	horseweed	ASTERACEAE	herb	3
<i>Coreopsis grandiflora</i>	no data	CORE GRAN	tickseed	ASTERACEAE	herb	5
<i>Cornus drummondii</i>	FAC	CORN DRUM	rough-leaved dogwood	CORNACEAE	shrub	3
<i>Corydalis crystallina</i>	no data	CORY CRYC	mealy fumewort	FUMARIACEAE	herb	5
<i>Crataegus mollis</i>	FAC	CRAT MOLL	hairy hawthorn	ROSACEAE	herb	8
<i>Crataegus reverchonii</i>	no data	CRAT CRUS	Reverchon's hawthorn	ROSACEAE	shrub	21
<i>Crotalaria sagittalis</i>	no data	CROT SAGI	rattlebox	FABACEAE	herb	12
<i>Croton capitatus</i>	no data	CROT CAPI	goatweed	EUPHORBIACEAE	herb	3
<i>Croton glandulosus</i> var. <i>septentrionalis</i>	no data	CROT GLAN SEPT	tropic croton	EUPHORBIACEAE	herb	3
<i>Croton monanthogynus</i>	no data	CROT MONA	prairie tea	EUPHORBIACEAE	herb	8
<i>Croton willdenowii</i>	no data	CROT WILD	rushfoil	EUPHORBIACEAE	herb	3
<i>Cruciata pedemontana</i> *	no data	CRUC PEDE	yellow-flowered bedstraw	RUBIACEAE	herb	5
<i>Cuscuta campestris</i>	no data	CUSC CAMP	field dodder	CONVOLVULACEAE	herb	12
<i>Cynodon dactylon</i> **	FACU	CYNO DACT	Bermuda grass	POACEAE	herb	3
<i>Cyperus acuminatus</i>	OBL	CYPE ACUM	tapertip flatsedge	CYPERACEAE	herb	6
<i>Cyperus echinatus</i>	FAC	CYPE ECHI	globe flatsedge	CYPERACEAE	herb	3
<i>Cyperus erythrorhizos</i>	OBL	CYPE ERYT	redroot flatsedge	CYPERACEAE	herb	11
<i>Cyperus esculentus</i>	FAC	CYPE ESCU	yellow nutsedge	CYPERACEAE	herb	3
<i>Cyperus flavescens</i>	OBL	CYPE FLAV	yellow flatsedge	CYPERACEAE	herb	3
<i>Cyperus lancastris</i>	FAC	CYPE LANC	manyflower flatsedge	CYPERACEAE	herb	21
<i>Cyperus lupulinus</i>	no data	CYPE LUPU	flatsedge	CYPERACEAE	herb	4
<i>Cyperus odoratus</i>	FACW	CYPE ODOR	rusty flatsedge	CYPERACEAE	herb	6
<i>Cyperus pseudovegetus</i>	FACW	CYPE PSEU	marsh flatsedge	CYPERACEAE	herb	3
<i>Cyperus strigosus</i>	FACW	CYPE STRI	false nutsedge	CYPERACEAE	herb	1
<i>Dactylis glomerata</i> **	FACU	DACT GLOM	orchard grass	POACEAE	herb	3
<i>Datura stramonium</i> *	no data	DATU STRA	Jimson weed	SOLANACEAE	herb	5
<i>Daucus carota</i> **	no data	DAUC CARO	Queen Anne's lace	APIACEAE	herb	3
<i>Desmanthus illinoensis</i>	FAC	DESM ILLI	Illinois bundleflower	FABACEAE	herb	21
<i>Desmodium canescens</i>	no data	DESM CANE	tick-trefoil	FABACEAE	herb	19
<i>Desmodium nuttallii</i>	no data	DESM NUTT	tick-trefoil	FABACEAE	herb	8
<i>Desmodium obtusum</i>	no data	DESM OBTU	tick-trefoil	FABACEAE	herb	8
<i>Desmodium paniculatum</i>	FACU	DESM PANI	tick-trefoil	FABACEAE	herb	3
<i>Desmodium sessilifolium</i>	no data	DESM SESS	sessile-leaf tick-trefoil	FABACEAE	herb	16
<i>Dianthus armeria</i> *	UPL	DIAN ARME	Deptford pink	CARYOPHYLLACEAE	herb	31
<i>Dichanthelium aciculare</i>	FACU	DICH ACIC	slimleaf rosettegrass	POACEAE	herb	3
<i>Dichanthelium acuminatum</i>	FAC	DICH ACUM	pointed rosettegrass	POACEAE	herb	3
<i>Dichanthelium clandestinum</i>	FACW	DICH CLAN	deer-tongue rosettegrass	POACEAE	herb	8

<i>Dichanthelium commutatum</i>	FAC	DICH COMM	variable rosettegrass	POACEAE	herb	11
<i>Dichanthelium dichotomum</i>	FAC	DICH DICH	rosettegrass	POACEAE	herb	3
<i>Dichanthelium malacophyllum</i>	no data	DICH MALA	soft-leaved rosettegrass	POACEAE	herb	19
<i>Dichanthelium oligosanthes</i> var. <i>scribnerianum</i>	FACU	DICH OLIG SCRI	Scribner's rosettegrass	POACEAE	herb	5
<i>Dichanthelium scoparium</i>	FACW	DICH SCOP	velvet rosettegrass	POACEAE	herb	3
<i>Dichanthelium sphaerocarpon</i>	FACU	DICH SPHA	rosettegrass	POACEAE	herb	5
<i>Digitaria ciliaris</i> **	FAC	DIGI CILI	southern crabgrass	POACEAE	herb	4
<i>Digitaria ischaemum</i> **	UPL	DIGI ISCH	smooth crabgrass	POACEAE	herb	3
<i>Diodia teres</i>	FACU-	DIOD TERE	poorjoe	RUBIACEAE	herb	3
<i>Diodia virginiana</i>	FACW	DIOD VIRG	Virginia buttonweed	RUBIACEAE	herb	3
<i>Diospyros virginiana</i>	FAC	DIOS VIRG	persimmon	EBENACEAE	tree/sapling	3
<i>Dysphania ambrosioides</i> *	FACU	DYSP AMBR	wormseed	CHENOPODIACEAE	herb	3
<i>Echinochloa colona</i> *	FACW	ECHI COLO	jungle rice	POACEAE	herb	3
<i>Echinochloa crus-galli</i> *	FACW-	ECHI CRUS	barnyard grass	POACEAE	herb	1
<i>Echinochloa muricata</i>	FAC	ECHI MURI	barnyard grass	POACEAE	herb	3
<i>Eclipta prostrata</i>	FACW-	ECLI PROS	yerba de tajo	ASTERACEAE	herb	1
<i>Eleocharis acicularis</i>	OBL	ELEO ACIC	least spikerush	CYPERACEAE	herb	5
<i>Eleocharis lanceolata</i>	FACW	ELEO LANC	spikerush	CYPERACEAE	herb	3
<i>Eleocharis macrostachya</i>	OBL	ELEO MACR	pale spikerush	CYPERACEAE	herb	16
<i>Eleocharis obtusa</i>	OBL	ELEO OBTU	blunt spikerush	CYPERACEAE	herb	3
<i>Eleocharis palustris</i>	OBL	ELEO PALU	common spikerush	CYPERACEAE	herb	3
<i>Eleocharis quadrangulata</i>	OBL	ELEO QUAD	squarestem spikerush	CYPERACEAE	herb	9
<i>Eleocharis tenuis</i> var. <i>verrucosa</i>	FACW	ELEO TENU VERR	slender spikerush	CYPERACEAE	herb	5
<i>Eleocharis wolfii</i>	OBL	ELEO WOLF	Wolf's spikerush	CYPERACEAE	herb	5
<i>Eleusine indica</i> *	FACU	ELEU INDI	India goosegrass	POACEAE	herb	3
<i>Elymus glabrifloris</i>	no data	ELYM GLAB	wild rye	POACEAE	herb	3
<i>Eragrostis hirsuta</i>	UPL	ERAG HIRS	bigtop lovegrass	POACEAE	herb	22
<i>Eragrostis spectabilis</i>	FACU	ERAG SPEC	purple lovegrass	POACEAE	herb	3
<i>Eragrostis intermedia</i>	no data	ERAG INTE	lovegrass	POACEAE	herb	8
<i>Erechtites hieraciifolia</i>	FAC-	EREC HIER	fireweed	ASTERACEAE	herb	8
<i>Erigeron annuus</i>	FACU	ERIG ANNU	fleabane	ASTERACEAE	herb	3
<i>Erigeron philadelphicus</i>	FACU	ERIG PHIL	Philadelphia fleabane	ASTERACEAE	herb	32
<i>Erigeron strigosus</i>	FAC	ERIG STRI	daisy fleabane	ASTERACEAE	herb	5
<i>Erigeron tenuis</i>	no data	ERIG TENU	fleabane	ASTERACEAE	herb	25
<i>Eryngium yuccifolium</i> +	FAC	ERYN YUCC	rattlensnake master	APIACEAE	herb	10
<i>Euonymus fortunei</i> **	no data	EUON FORT	winter-creeper	CELASTRACEAE	woody vine	19
<i>Eupatorium perfoliatum</i>	FACW+	EUPA PERF	clasping boneset	ASTERACEAE	herb	3
<i>Eupatorium serotinum</i>	FAC	EUPA SERO	late boneset	ASTERACEAE	herb	1
<i>Euphorbia spathulata</i>	FACU	EUPH SPAT	warty spurge	EUPHORBIACEAE	herb	21
<i>Euthamia gymnospermoides</i>	FAC	EUTH GYMN	Texas goldentop	ASTERACEAE	herb	21
<i>Festuca rubra</i>	FACU+	FEST RUBR	red fescue	POACEAE	herb	2
<i>Fimbristylis annua</i>	FACW	FIMB ANNU	annual fimbry	CYPERACEAE	herb	9
<i>Fimbristylis puberula</i>	OBL	FIMB PUBE	hairy fimbry	CYPERACEAE	herb	5
<i>Fleischmannia incarnata</i>	FAC	FLEI INCA	pink thoroughwort	ASTERACEAE	herb	31
<i>Fraxinus pennsylvanica</i>	FACW	FRAX PENN	green ash	OLEACEAE	tree/sapling	3
<i>Galactia regularis</i>	no data	GALA REGU	milk pea	FABACEAE	herb	3
<i>Galium aparine</i>	FACU	GALI APAR	cleavers	RUBIACEAE	herb	19
<i>Galium obtusum</i>	FACW-	GALI OBTU	bluntleaf bedstraw	RUBIACEAE	herb	7
<i>Galium pilosum</i>	no data	GALI PILO	hairy bedstraw	RUBIACEAE	herb	4
<i>Gamochaeta antillana</i>	no data	GAMO ANTI	cudweed	ASTERACEAE	herb	19
<i>Gamochaeta purpurea</i>	UPL	GAMO PURP	purple cudweed	ASTERACEAE	herb	5

<i>Gaura longiflora</i>	no data	GAUR LONG	gaura	ONAGRACEAE	herb	8
<i>Geranium carolinianum</i>	no data	GERA CARO	Carolina cranesbill	GERANIACEAE	herb	5
<i>Geranium dissectum</i> *	no data	GERA DISS	cutleaf cranesbill	GERANIACEAE	herb	5
<i>Geranium molle</i> *	no data	GERA MOLL	dovesfoot cranesbill	GERANIACEAE	herb	4
<i>Geum canadense</i>	FACU	GEUM CANA	white avens	ROSACEAE	herb	19
<i>Glandularia canadensis</i>	no data	GLAN CANA	rose vervain	VERBENACEAE	herb	3
<i>Gleditsia triacanthos</i>	FAC-	GLED TRIA	honey locust	FABACEAE	tree/sapling	3
<i>Glyceria septentrionalis</i>	OBL	GLYC SEPT	mannagrass	POACEAE	herb	3
<i>Gratiola neglecta</i>	OBL	GRAT NEGL	hedge-hyssop	SCROPHULARIACEAE	herb	9
<i>Gratiola virginiana</i>	OBL	GRAT VIRG	hedge-hyssop	SCROPHULARIACEAE	herb	3
<i>Helenium amarum</i>	FACU-	HELE AMAR	bitterweed	ASTERACEAE	herb	3
<i>Helenium flexuosum</i>	FACW	HELE FLEX	purple-headed sneezeweed	ASTERACEAE	herb	1
<i>Helenium autumnale</i>	FACW	HELE AUTU	fall sneezeweed	ASTERACEAE	herb	8
<i>Helianthus grosseserratus</i>	FAC+	HELI GROS	sawtooth sunflower	ASTERACEAE	herb	3
<i>Helianthus mollis</i>	no data	HELI MOLL	ashy sunflower	ASTERACEAE	herb	3
<i>Heliotropium indicum</i> *	FAC	HELI INDI	Indian heliotrope	BORAGINACEAE	herb	21
<i>Hibiscus moscheutos</i> ssp. <i>lasiocarpus</i>	OBL	HIBI MOSC LASI	rose mallow	MALVACEAE	herb	3
<i>Hieracium gronovii</i>	UPL	HIER GRON	hawkweed	ASTERACEAE	herb	2
<i>Hordeum pusillum</i> *	FACU	HORD PUSI	little barley	POACEAE	herb	5
<i>Hypericum drummondii</i>	FACU	HYPE DRUM	nits-and-lice	CLUSIACEAE	herb	8
<i>Hypericum gymnanthum</i>	FACW	HYPE GYMN	clasping St. John's wort	CLUSIACEAE	herb	10
<i>Hypericum hypericoides</i> var. <i>multicaule</i>	FAC	HYPE HYPE MULT	creeping St. Andrew's cross	CLUSIACEAE	shrub	3
<i>Hypericum mutilum</i>	FACW	HYPE MUTI	dwarf St. John's wort	CLUSIACEAE	herb	9
<i>Hypericum punctatum</i>	FAC	HYPE PUNC	dotted St. John's wort	CLUSIACEAE	herb	8
<i>Ilex decidua</i>	FACW	ILEX DECI	deciduous holly	AQUIFOLIACEAE	shrub	22
<i>Ipomoea lacunosa</i>	FAC+	IPOM LACU	whitestar morning glory	CONVOLVULACEAE	herb	10
<i>Ipomoea pandurata</i>	FACU	IPOM PAND	wild potato vine	CONVOLVULACEAE	herb	5
<i>Isoetes melanopoda</i>	OBL	ISOE MELA	black-footed quillwort	ISOETACEAE	herb	19
<i>Isolepis carinata</i>	FACW+	ISOL CARI	bulrush	CYPERACEAE	herb	5
<i>Iva annua</i>	FAC	IVA ANNU	sumpweed	ASTERACEAE	herb	32
<i>Juncus anthelatus</i>	no data	JUNC ANTH	rush	JUNCACEAE	herb	3
<i>Juncus biflorus</i>	FACW	JUNC BIFL	rush	JUNCACEAE	herb	3
<i>Juncus brachycarpus</i>	FACW	JUNC BRAC	rush	JUNCACEAE	herb	9
<i>Juncus diffusissimus</i>	FACW	JUNC DIFF	spreading rush	JUNCACEAE	herb	10
<i>Juncus effusus</i>	FACW+	JUNC EFFU	soft rush	JUNCACEAE	herb	3
<i>Juncus interior</i>	FACU	JUNC INTE	inland rush	JUNCACEAE	herb	5
<i>Juncus marginatus</i>	FACW	JUNC MARG	rush	JUNCACEAE	herb	8
<i>Juncus secundus</i>	FAC	JUNC SECU	rush	JUNCACEAE	herb	4
<i>Juncus tenuis</i>	FAC	JUNC TENU	path rush	JUNCACEAE	herb	24
<i>Juncus torreyi</i>	FACW	JUNC TORR	Torrey's rush	JUNCACEAE	herb	23
<i>Juncus validus</i>	FACW+	JUNC VALI	rush	JUNCACEAE	herb	5
<i>Juniperus virginiana</i>	FACU-	JUNI VIRG	eastern redcedar	CUPRESSACEAE	tree/sapling	8
<i>Krigia dandelion</i>	FACU	KRIG DAND	potato dandelion	ASTERACEAE	herb	7
<i>Kummerowia stipulacea</i> **	FACU-	KUMM STIP	Korean bushclover	FABACEAE	herb	3
<i>Kummerowia striata</i> **	FACU	KUMM STRI	Japanese bushclover	FABACEAE	herb	3
<i>Lactuca canadensis</i>	FACU-	LACT CANA	Canada wild lettuce	ASTERACEAE	herb	16
<i>Lactuca saligna</i> *	UPL	LACT SALI	willowleaf lettuce	ASTERACEAE	herb	21
<i>Lactuca serriola</i> *	FAC	LACT SERR	prickly wild lettuce	ASTERACEAE	herb	3
<i>Leersia oryzoides</i>	OBL	LEER ORYZ	rice cutgrass	POACEAE	herb	2
<i>Leersia virginica</i>	FACW	LEER VIRG	Virginia cutgrass	POACEAE	herb	8

<i>Lemna minuta</i>	OBL	LEMN MINU	duckweed	LEMNACEAE	herb	5
<i>Lepidium virginicum</i>	FACU	LEPI VIRG	Virginia peppergrass	BRASSICACEAE	herb	3
<i>Lespedeza cuneata</i> **	NI	LESP CUNE	sericea lespedeza	FABACEAE	herb	3
<i>Lespedeza repens</i>	no data	LESP REPE	creeping lespedeza	FABACEAE	herb	21
<i>Leucospora multifida</i>	OBL	LEUC MULT	leucospora	SCROPHULARIACEAE	herb	5
<i>Liatris pycnostachya</i> +	FACU	LIAT PYCN	prairie gayfeather	ASTERACEAE	herb	27
<i>Liatris squarrulosa</i>	FACU	LIAT SQUA	Appalachian Blazing Star	ASTERACEAE	herb	29
<i>Ligustrum sinense</i> **	FAC	LIGU SINE	Chinese privet	OLEACEAE	shrub	17
<i>Lindernia dubia</i> var. <i>anagallidea</i>	OBL	LIND ANAG	false pimpernel	SCROPHULARIACEAE	herb	10
<i>Lindernia dubia</i> var. <i>dubia</i>	OBL	LIND DUBI	false pimpernel	SCROPHULARIACEAE	herb	23
<i>Linum medium</i> var. <i>texanum</i>	FACU	LINU MEDI TEXA	stiff yellow flax	LINACEAE	herb	23
<i>Lobelia siphilitica</i>	OBL	LOBE SIPH	big blue lobelia	CAMPANULACEAE	herb	8
<i>Lobelia spicata</i>	FAC	LOBE SPIC	spike lobelia	CAMPANULACEAE	herb	5
<i>Lolium perenne</i> *	FACU	LOLI PERE	ryegrass	POACEAE	herb	5
<i>Lonicera japonica</i> **	FAC-	LONI JAPO	Japanese honeysuckle	CAPRIFOLIACEAE	woody vine	3
<i>Lonicera maackii</i> **	no data	LONI MAAC	bush honeysuckle	CAPRIFOLIACEAE	shrub	17
<i>Lonicera sempervirens</i>	FAC	LONI SEMP	trumpet honeysuckle	CAPRIFOLIACEAE	woody vine	5
<i>Ludwigia alternifolia</i>	OBL	LUDW ALTE	seedbox	ONAGRACEAE	herb	8
<i>Ludwigia glandulosa</i>	OBL	LUDW GLAN	primrose-willow	ONAGRACEAE	herb	23
<i>Ludwigia palustris</i>	OBL	LUDW PALU	creeping seedbox	ONAGRACEAE	herb	3
<i>Ludwigia peploides</i> ssp. <i>glabrescens</i>	OBL	LUDW PEPL GLAB	floating primrose-willow	ONAGRACEAE	herb	3
<i>Luzula echinata</i>	FAC	LUZU ECHI	wood rush	JUNCACEAE	herb	10
<i>Lycopus americanus</i>	OBL	LYCO AMER	American water horehound	LAMIACEAE	herb	8
<i>Lysimachia lanceolata</i>	FAC	LYSI LANC	lanceleaf loosestrife	PRIMULACEAE	herb	32
<i>Lythrum alatum</i>	FACW+	LYTH ALAT	winged loosestrife	LYTHRACEAE	herb	5
<i>Maclura pomifera</i> *	FACU	MACL POMI	bois d'arc	MORACEAE	tree/sapling	3
<i>Mecardonia acuminata</i>	FACW	MECA ACUM	purple axillflower	SCROPHULARIACEAE	herb	3
<i>Medicago lupulina</i>	no data	MEDI SP.	black medic	FABACEAE	herb	16
<i>Melilotus albus</i> **	FACU-	MELI ALBU	white sweetclover	FABACEAE	herb	3
<i>Melilotus officinalis</i> *	FACU-	MELI OFFI	yellow sweetclover	FABACEAE	herb	16
<i>Melothria pendula</i>	FACW-	MELO PEND	dwarf cucumber vine	CUCURBITACEAE	herb	10
<i>Microstegium vimineum</i> **	FAC	MICR VIMI	Japanese stiltgrass	POACEAE	herb	
<i>Mimosa quadrivalvis</i> var. <i>nuttallii</i>	no data	MIMO QUAD NUTT	sensitive brier	FABACEAE	herb	3
<i>Miscanthus sinensis</i> **	FACU	MISC SINE	Chinese Silvergrass	POACEAE	herb	29
<i>Mollugo verticillata</i>	FAC	MOLL VERT	green carpetweed	MOLLUGINACEAE	herb	10
<i>Morus alba</i> *	UPL	MORU ALBA	white mulberry	MORACEAE	tree/sapling	20
<i>Morus rubra</i>	FAC	MORU RUBR	red mulberry	MORACEAE	tree/sapling	8
<i>Muhlenbergia schreberi</i>	FAC	MUHL SCHR	nimblewill	POACEAE	herb	8
<i>Muhlenbergia</i> sp.	no data	MUHL SP.	muhly grass	POACEAE	herb	19
<i>Myosotis macrosperma</i>	FAC	MYOS MACR	large-seeded forget-me-not	BORAGINACEAE	herb	19
<i>Myriophyllum</i> sp.	OBL	MYRI SP	water milfoil	HALORAGACEAE	herb	9
<i>Nelumbo lutea</i>	OBL	NELU LUTE	American yellow lotus	Nelumbonaceae	herb	30
<i>Nothoscordum bivalve</i>	FAC	NOTH BIVA	crow poison	ALLIACEAE	herb	7
<i>Nuttallanthus texanus</i>	no data	NUTT TEXA	blue toadflax	SCROPHULARIACEAE	herb	5
<i>Oenothera biennis</i>	FACU	OENO BIEN	evening-primrose	ONAGRACEAE	herb	7
<i>Oenothera laciniata</i>	FACU	OENO LACI	cutleaf evening-primrose	ONAGRACEAE	herb	5
<i>Orbexilum pedunculatum</i> var. <i>pedunculatum</i>	FACU	ORBE PEDU	Sampson's snakeroot	FABACEAE	herb	5
<i>Oxalis dillenii</i>	no data	OXAL DILL	yellow wood sorrel	OXALIDACEAE	herb	3
<i>Oxalis stricta</i>	FACU	OXAL STRI	yellow wood sorrel	OXALIDACEAE	herb	27
<i>Oxalis violacea</i>	no data	OXAL VIOL	violet woodsorrel	OXALIDACEAE	herb	5

<i>Panicum anceps</i>	FAC-	PANI ANCE	beaked panicgrass	POACEAE	herb	3
<i>Panicum capillare</i>	FAC	PANI CAPI	witchgrass	POACEAE	herb	8
<i>Panicum dichotomiflorum</i>	FACW	PANI DICH	fall panicgrass	POACEAE	herb	3
<i>Panicum rigidulum</i>	FACW	PANI RIGI	rigid panicgrass	POACEAE	herb	5
<i>Panicum virgatum</i>	FAC+	PANI VIRG	switchgrass	POACEAE	herb	3
<i>Parthenocissus quinquefolia</i>	FACU	PART QUIN	Virginia creeper	VITACEAE	woody vine	19
<i>Paspalum dilatatum *</i>	FAC+	PASP DILA	Dallisgrass	POACEAE	herb	3
<i>Paspalum floridanum</i>	FACW-	PASP FLOR	Florida crowngrass	POACEAE	herb	3
<i>Paspalum laeve</i>	FACW-	PASP LAEV	field paspalum	POACEAE	herb	3
<i>Paspalum notatum *</i>	FACU+	PASP NOTA	Bahia grass	POACEAE	herb	10
<i>Paspalum pubiflorum</i>	FACW	PASP PUBI	hairyseed crowngrass	POACEAE	herb	10
<i>Paspalum setaceum</i>	FAC	PASP SETA	thin crowngrass	POACEAE	herb	4
<i>Passiflora incarnata</i>	no data	PASS INCA	passion flower	PASSIFLORACEAE	herb	3
<i>Passiflora lutea</i>	no data	PASS LUTE	yellow passion flower	PASSIFLORACEAE	herb	19
<i>Penstemon digitalis</i>	FAC	PENS DIGI	foxglove beard-tongue	SCROPHULARIACEAE	herb	5
<i>Penstemon tubaeiflorus</i>	no data	PENS TUBA	whitewand beard-tongue	SCROPHULARIACEAE	herb	3
<i>Penthorum sedoides</i>	OBL	PENT SEDO	ditch stonecrop	CRASSULACEAE	herb	31
<i>Perilla frutescens **</i>	FACU	PERI FRUT	beefsteak plant	LAMIACEAE	herb	32
<i>Persicaria hydropiper *</i>	OBL	PERS HYDROPIPER	water pepper	POLYGONACEAE	herb	6
<i>Persicaria hydropiperoides</i>	OBL	PERS HYDROPIPEROIDES	wild water pepper	POLYGONACEAE	herb	3
<i>Persicaria lapathifolia</i>	FACW	PERS LAPA	pale smartweed	POLYGONACEAE	herb	3
<i>Persicaria longiseta *</i>	no data	PERS LONG	pink smartweed	POLYGONACEAE	herb	8
<i>Persicaria maculosa *</i>	FACW	PERS MACU	lady's-thumb	POLYGONACEAE	herb	6
<i>Persicaria pensylvanica</i>	FACW	PERS PENS	Pennsylvania smartweed	POLYGONACEAE	herb	3
<i>Persicaria punctata</i>	FACW+	PERS PUNC	dotted smartweed	POLYGONACEAE	herb	2
<i>Phleum pratense *</i>	FACU	PHLE PRAT	timothy	POACEAE	herb	21
<i>Phyla lanceolata</i>	OBL	PHYL LANC	lanceleaf fogfruit	VERBENACEAE	herb	24
<i>Physalis angulata</i>	FAC	PHYS ANGU	smooth groundcherry	SOLANACEAE	herb	8
<i>Physalis heterophylla</i>	no data	PHYS HETE	clammy groundcherry	SOLANACEAE	herb	10
<i>Physalis longifolia</i>	no data	PHYS LONG	longleaf groundcherry	SOLANACEAE	herb	10
<i>Physalis pubescens</i>	FACU	PHYS PUBE	hairy groundcherry	SOLANACEAE	herb	3
<i>Physostegia angustifolia</i>	FACW	PHYS ANGU	false dragonhead	LAMIACEAE	herb	3
<i>Phytolacca americana</i>	FACU+	PHYT AMER	pokeweed	PHYTOLACACEAE	herb	3
<i>Plantago aristata</i>	no data	PLAN ARIS	bracted plantain	PLANTAGINACEAE	herb	3
<i>Plantago lanceolata *</i>	FAC	PLAN LANC	English plantain	PLANTAGINACEAE	herb	3
<i>Plantago rugelii</i>	FAC	PLAN RUGE	blackseed plantain	PLANTAGINACEAE	herb	2
<i>Plantago virginica</i>	FACU-	PLAN VIRG	Virginia plantain	PLANTAGINACEAE	herb	5
<i>Platanus occidentalis</i>	FACW-	PLAT OCCI	American sycamore	PLATANACEAE	tree/sapling	12
<i>Pluchea camphorata</i>	FACW	PLUC CAMP	stinkweed	ASTERACEAE	herb	27
<i>Poa annua *</i>	FAC	POA ANNU	annual bluegrass	POACEAE	herb	5
<i>Poa compressa *</i>	FACU-	POA COMP	Canada bluegrass	POACEAE	herb	3
<i>Poa pratensis *</i>	FACU+	POA PRAT	Kentucky bluegrass	POACEAE	herb	5
<i>Polygala incarnata</i>	FAC-	POLY INCA	pink milkwort	POLYGALACEAE	herb	16
<i>Polygala sanguinea</i>	FAC-	POLY SANG	purple milkwort	POLYGALACEAE	herb	10
<i>Polygala verticillata</i>	UPL	POLY VERT	whorled milkwort	POLYGALACEAE	herb	23
<i>Polygonum aviculare *</i>	FAC-	POLY AVIC	knotweed	POLYGONACEAE	herb	3
<i>Polygonum erectum</i>	FACU	POLY EREC	erect knotweed	POLYGONACEAE	herb	10
<i>Populus deltoides</i>	FAC+	POPU DELT	eastern cottonwood	SALICACEAE	tree/sapling	10
<i>Potamogeton diversifolius</i>	OBL	POTA DIVE	pondweed	POTAMOGETONACEAE	herb	5
<i>Potamogeton nodosus</i>	OBL	POTA NODO	pondweed	POTAMOGETONACEAE	herb	1
<i>Potamogeton pusillus</i>	OBL	POTA PUSI	narrowleaf pondweed	POTAMOGETONACEAE	herb	11
<i>Potentilla recta *</i>	no data	POTE RECT	rough-fruited cinquefoil	ROSACEAE	herb	5

<i>Potentilla simplex</i>	FACU	POTE SIMP	cinquefoil	ROSACEAE	herb	5
<i>Proserpinaca palustris</i>	OBL	PROS PALU	mermaid weed	HALORAGACEAE	herb	1
<i>Prunella vulgaris</i> ssp. lanceolata	FAC-	PRUN VULG	heal-all	LAMIACEAE	herb	4
<i>Prunus munsoniana</i>	no data	PRUN MUNS	wild goose plum	ROSACEAE	shrub	19
<i>Prunus serotina</i>	FACU	PRUN SERO	black cherry	ROSACEAE	tree/sapling	3
<i>Pseudognaphalium obtusifolium</i>	no data	PSEU OBTU	rabbit-tobacco	ASTERACEAE	herb	22
<i>Pycnanthemum pilosum</i>	UPL	PYCN PILO	hairy mountain mint	LAMIACEAE	herb	7
<i>Pycnanthemum tenuifolium</i>	FAC-	PYCN TENU	slender mountain mint	LAMIACEAE	herb	3
<i>Pycnanthemum pilosum</i> X <i>P. tenuifolium</i>	no data	PYCN PILO X TENU	hybrid mountain mint	LAMIACEAE	herb	13
<i>Pyrrhopappus carolinianus</i>	no data	PYRR CARO	false dandelion	ASTERACEAE	herb	12
<i>Pyrus calleryana</i> **	no data	PYRU CALL	callery pear	ROSACEAE	tree/sapling	3
<i>Quercus</i> +	-	QUER SP.	oak	FAGACEAE	tree/sapling	10
<i>Ranunculus bulbosus</i> *	FAC+	RANU BULB	bulbous buttercup	RANUNCULACEAE	herb	1
<i>Ranunculus laxicaulis</i>	OBL	RANU LAXI	water plantain spearwort	RANUNCULACEAE	herb	5
<i>Ranunculus micranthus</i>	FACU	RANU MICR	rock buttercup	RANUNCULACEAE	herb	19
<i>Ranunculus parviflorus</i> *	FAC	RANU PARV	smallflower crowfoot	RANUNCULACEAE	herb	5
<i>Ranunculus sardous</i> *	FAC+	RANU SARD	hairy buttercup	RANUNCULACEAE	herb	3
<i>Rhexia mariana</i>	FACW+	RHEX MARI	meadow beauty	MELASTOMACEAE	herb	10
<i>Rhus copallinum</i>	NI	RHUS COPA	winged sumac	ANACARDIACEAE	shrub	10
<i>Rhus glabra</i>	no data	RHUS GLAB	smooth sumac	ANACARDIACEAE	shrub	10
<i>Rhynchospora harveyi</i>	OBL	RHYN HARV	Harvey's beaksedge	CYPERACEAE	herb	5
<i>Rhynchospora macrostachya</i>	OBL	RHYN MACR	tall horned beaksedge	CYPERACEAE	herb	1
<i>Rhynchospora recognita</i>	FACW	RHYN RECO	beaksedge	CYPERACEAE	herb	14
<i>Rorippa palustris</i> ssp. fernaldiana	OBL	RORI PALU FERN	Fernald's yellowcress	BRASSICACEAE	herb	12
<i>Rosa carolina</i>	FACU	ROSA CARO	Carolina rose	ROSACEAE	shrub	3
<i>Rosa multiflora</i> **	UPL	ROSA MULT	multiflora rose	ROSACEAE	shrub	3
<i>Rosa setigera</i>	FACU	ROSA SETI	prairie rose	ROSACEAE	shrub	3
<i>Rotala ramosior</i>	OBL	ROTA RAMO	toothcup	LYTHRACEAE	herb	3
<i>Rubus aboriginum</i>	no data	RUBU ABOR	dewberry	ROSACEAE	shrub	21
<i>Rubus laudatus</i>	no data	RUBU LAUD	plains blackberry	ROSACEAE	shrub	21
<i>Rubus satis</i>	no data	RUBU SATI	dewberry	ROSACEAE	shrub	21
<i>Rubus serissimus</i> **	UPL	RUBU SERI	Himalayan blackberry	ROSACEAE	shrub	5
<i>Rubus flagellaris</i>	UPL	RUBU FLAG	northern dewberry	ROSACEAE	herb	3
<i>Rudbeckia hirta</i>	FACU	RUDB HIRT	black-eyed Susan	ASTERACEAE	herb	3
<i>Rudbeckia subtomentosa</i>	FAC+	RUDB SUBT	sweet coneflower	ASTERACEAE	herb	3
<i>Ruellia humilis</i> var. <i>humilis</i>	FACU	RUEL HUMI	hairy wild petunia	ACANTHACEAE	herb	3
<i>Rumex acetosella</i> *	FACU+	RUME ACET	red sorrel	POLYGONACEAE	herb	5
<i>Rumex altissimus</i>	FACW	RUME ALTI	pale dock	POLYGONACEAE	herb	4
<i>Rumex crispus</i> **	FAC	RUME CRIS	curly dock	POLYGONACEAE	herb	3
<i>Rumex obtusifolius</i> **	FACU	RUME OBTU	bitter dock	POLYGONACEAE	herb	23
<i>Sabatia angularis</i>	FAC	SABA ANGU	winged rosepink	GENTIANACEAE	herb	5
<i>Sabatia campestris</i>	FACU	SABA CAMP	prairie rosepink	GENTIANACEAE	herb	7
<i>Sagittaria montevidensis</i>	OBL	SAGI MONT	duck potato	ALISMACEAE	herb	3
<i>Sagittaria platyphylla</i>	OBL	SAGI PLAT	delta arrowhead	ALISMACEAE	herb	23
<i>Salix nigra</i>	OBL	SALI NIGR	black willow	SALICACEAE	tree/sapling	3
<i>Salsola tragus</i> *	FACU	SALS TRAG	Russian thistle	CHENOPODIACEAE	herb	11
<i>Salvia lyrata</i>	FAC-	SALV LYRA	cancerweed	LAMIACEAE	herb	3
<i>Sambucus nigra</i> ssp. <i>canadensis</i>	FAC	SAMB NIGR CANA	elderberry	CAPRIFOLIACEAE	shrub	24
<i>Sanicula canadensis</i>	UPL	SANI CANA	Canada black snakeroot	APIACEAE	herb	19
<i>Sassafras albidum</i>	FACU	SASS ALBI	sassafras	LAURACEAE	tree/sapling	3
<i>Schedonorus arundinaceus</i> **	FAC-	SCHE ARUN	tall fescue	POACEAE	herb	3

<i>Schizachyrium scoparium</i>	FACU	SCHI SCOP	little bluestem	POACEAE	herb	3
<i>Schoenoplectus tabernaemontani</i>	OBL	SCHO TABE	softstem bulrush	CYPERACEAE	herb	9
<i>Scirpus cyperinus</i>	FACW	SCIR CYPE	woolgrass bulrush	CYPERACEAE	herb	20
<i>Scirpus georgianus</i>	OBL	SCIR GEOR	Georgia bulrush	CYPERACEAE	herb	3
<i>Scirpus pendulus</i>	OBL	SCIR PEND	drooping bulrush	CYPERACEAE	herb	5
<i>Scleria ciliata</i>	FAC	SCLE CILI	fringed nutrush	CYPERACEAE	herb	12
<i>Scleria pauciflora</i> var. <i>caroliniana</i>	FAC+	SCLE PAUC	fewflower nutrush	CYPERACEAE	herb	5
<i>Scutellaria parvula</i> var. <i>missouriensis</i>	no data	SCUT PARV MISS	glade skullcap	LAMIACEAE	herb	27
<i>Senna marilandica</i>	FACW	SENN MARI	wild senna	FABACEAE	herb	30
<i>Setaria faberi</i> *	UPL	SETA FABE	Chinese foxtail	POACEAE	herb	3
<i>Setaria italica</i> *	FACU	SETA ITAL	Italian foxtail	POACEAE	herb	14
<i>Setaria parviflora</i>	FAC	SETA PARV	knotroot bristlegrass	POACEAE	herb	3
<i>Setaria pumila</i> ssp. <i>pumila</i> *	FAC	SETA PUMI	yellow foxtail	POACEAE	herb	3
<i>Setaria viridis</i> *	no data	SETA VIRI	green bristlegrass	POACEAE	herb	23
<i>Sherardia arvensis</i> *	no data	SHER ARVE	field madder	RUBIACEAE	herb	5
<i>Sida spinosa</i> *	FACU	SIDA SPIN	prickly sida	MALVACEAE	herb	3
<i>Sideroxylon lanuginosum</i>	FACU	SIDE LANU	chittum wood	SAPOTACEAE	tree/sapling	3
<i>Silene antirrhina</i>	no data	SILE ANTI	sleepy catchfly	CARYOPHYLLACEAE	herb	19
<i>Silphium laciniatum</i>	no data	SILP LACI	compass plant	ASTERACEAE	herb	7
<i>Sisymbrium officinale</i> *	no data	SISY OFFI	hedge mustard	BRASSICACEAE	herb	5
<i>Sisyrinchium angustifolium</i>	FAC	SISY ANGU	blue-eyed grass	IRIDACEAE	herb	7
<i>Sisyrinchium atlanticum</i>	FACW-	SISY ATLA	blue-eyed grass	IRIDACEAE	herb	5
<i>Smallanthus uvedalia</i>	UPL	SMAL UVED	hairy leafcup	ASTERACEAE	herb	31
<i>Smilax bona-nox</i>	FAC	SMIL BONA	bull greenbrier	SMILACACEAE	woody vine	3
<i>Smilax rotundifolia</i>	FAC	SMIL ROTU	common greenbrier	SMILACACEAE	woody vine	19
<i>Solanum carolinense</i>	FACU	SOLA CARO	Carolina horsenettle	SOLANACEAE	herb	3
<i>Solanum sarrachoides</i> *	no data	SOLA PHYS	hairy nightshade	SOLANACEAE	herb	6
<i>Solidago altissima</i>	FACU	SOLI ALTI	tall goldenrod	ASTERACEAE	herb	3
<i>Solidago gigantea</i>	FACW	SOLI GIGA	giant goldenrod	ASTERACEAE	herb	16
<i>Solidago rugosa</i>	FAC	SOLI RUGO	wrinkleleaf goldenrod	ASTERACEAE	herb	13
<i>Sonchus asper</i> **	FAC+	SONC ASPE	spiny sowthistle	ASTERACEAE	herb	10
<i>Sorghastrum nutans</i>	FACU	SORG NUTA	Indiangrass	POACEAE	herb	3
<i>Sorghum bicolor</i> *	FACU	SORG BICO	sorghum	POACEAE	herb	15
<i>Sorghum halepense</i> **	FACU	SORG HALE	Johnsongrass	POACEAE	herb	3
<i>Sphenopholis obtusata</i>	FAC+	SPHE OBTU	prairie wedgescale	POACEAE	herb	3
<i>Spiranthes cernua</i>	FACW	SPIR CERN	nodding ladies'-tresses	ORCHIDACEAE	herb	4
<i>Spiranthes vernalis</i>	FACW-	SPIR VERN	spring ladies'-tresses	ORCHIDACEAE	herb	10
<i>Spirodella polyrhiza</i>	OBL	SPIR POLY	giant duckweed	LEMNACEAE	herb	5
<i>Sporobolus compositus</i> var. <i>compositus</i>	UPL	SPOR COMP COMP	rough dropseed	POACEAE	herb	4
<i>Sporobolus compositus</i> var. <i>macer</i>	no data	SPOR COMP MACE	creeping dropseed	POACEAE	herb	22
<i>Sporobolus vaginiflorus</i> var. <i>vaginiflorus</i>	UPL	SPOR VAGI	dropseed	POACEAE	herb	8
<i>Steinchisma hians</i>	OBL	STEI HIAN	gaping panicgrass	POACEAE	herb	1
<i>Stellaria media</i> *	FACU	STEL MEDI	common chickweed	CARYOPHYLLACEAE	herb	5
<i>Strophostyles leiosperma</i>	no data	STRO LEIO	wild bean	FABACEAE	herb	8
<i>Strophostyles helvola</i>	FAC	STRO HELV	amberique-bean	FABACEAE	herb	3
<i>Stylosanthes biflora</i>	no data	STYL BIFL	pencil flower	FABACEAE	herb	7
<i>Symphoricarpos orbiculatus</i>	FAC-	SYMP ORBI	coralberry	CAPRIFOLIACEAE	shrub	3
<i>Symphyotrichum divaricatum</i>	OBL	SYMP DIVA	annual aster	ASTERACEAE	herb	24
<i>Symphyotrichum dumosum</i>	FAC	SYMP DUMO	aster	ASTERACEAE	herb	1

<i>Symphytotrichum ericoides</i>	UPL	SYMP ERIC	heath aster	ASTERACEAE	herb	3
<i>Symphytotrichum lanceolatum</i>	NI	SYMP LANC	tall white aster	ASTERACEAE	herb	5
<i>Symphytotrichum patens</i>	no data	SYMP PATE	spreading aster	ASTERACEAE	herb	4
<i>Symphytotrichum pilosum</i>	FAC-	SYMP PILO	white heath aster	ASTERACEAE	herb	3
<i>Taraxacum officinale</i>	FACU	TARA OFFI	common dandelion	ASTERACEAE	herb	3
<i>Teucrium canadense</i>	FACW-	TEUC CANA	germander	LAMIACEAE	herb	10
<i>Torilis arvensis</i> *	no data	TORI ARVE	hedge parsley	APIACEAE	herb	3
<i>Toxicodendron radicans</i>	FAC	TOXI RAD	poison ivy	ANACARDIACEAE	woody vine	3
<i>Trachelospermum difforme</i>	FACW	TRAC DIFF	climbing dogbane	APOCYNACEAE	woody vine	5
<i>Tragia ramosa</i>	no data	TRAG RAMO	noseburn	EUPHORBIACEAE	herb	5
<i>Tridens flavus</i> var. <i>flavus</i>	FACU	TRID FLAV	purpletop tridens	POACEAE	herb	1
<i>Tridens strictus</i>	FACW	TRID STRI	longspike tridens	POACEAE	herb	1
<i>Tridens X oklahomensis</i>	no data	TRID OKLA	Oklahoma purpletop	POACEAE	herb	11
<i>Trifolium campestre</i> *	no data	TRIF CAMP	hop clover	FABACEAE	herb	5
<i>Trifolium dubium</i> *	FACU-	TRIF DUBI	low hop clover	FABACEAE	herb	5
<i>Trifolium pratense</i> *	FACU-	TRIF PRAT	red clover	FABACEAE	herb	3
<i>Trifolium repens</i> *	FACU	TRIF REPE	white clover	FABACEAE	herb	2
<i>Triodanis perfoliata</i> var. <i>biflora</i>	no data	TRIO PERF BIFL	round-leaved Venus' looking glass	CAMPANULACEAE	herb	19
<i>Triodanis perfoliata</i> var. <i>perfoliata</i>	FAC	TRIO PERF PERF	twinflower Venus' looking glass	CAMPANULACEAE	herb	5
<i>Tripsacum dactyloides</i> +	FACW	TRIP DACT	eastern gamagrass	POACEAE	herb	19
<i>Typha angustifolia</i> **	OBL	TYPH ANGU	narrowleaf cattail	TYPHACEAE	herb	7
<i>Typha domingensis</i>	OBL	TYPH DOMI	southern cattail	TYPHACEAE	herb	3
<i>Typha latifolia</i>	OBL	TYPH LATI	broadleaf cattail	TYPHACEAE	herb	14
<i>Ulmus alata</i>	FACU+	ULMU ALAT	winged elm	ULMACEAE	tree/sapling	3
<i>Ulmus americana</i>	FACW	ULMU AMER	American elm	ULMACEAE	tree/sapling	3
<i>Valerianaella radiata</i>	FAC	VALE RAD	cornsalad	VALERIANACEAE	herb	5
<i>Verbascum thapsus</i> *	no data	VERB THAP	woolly mullein	SCROPHULARIACEAE	herb	11
<i>Verbena bracteata</i>	FACU-	VERB BRAC	bigbract vervain	VERBENACEAE	herb	16
<i>Verbena hastata</i>	FAC	VERB HAST	blue vervain	VERBENACEAE	herb	3
<i>Verbena simplex</i>	OBL	VERB SIMP	vervain	VERBENACEAE	herb	5
<i>Verbena stricta</i>	no data	VERB STRI	hoary vervain	VERBENACEAE	herb	21
<i>Verbena urticifolia</i>	FAC+	VERB URTI	white vervain	VERBENACEAE	herb	5
<i>Vernonia arkansana</i>	FAC	VERN ARKA	Arkansas ironweed	ASTERACEAE	herb	10
<i>Vernonia baldwinii</i>	UPL	VERN BALD	Baldwin's ironweed	ASTERACEAE	herb	8
<i>Vernonia missurica</i>	FAC+	VERN MISS	Missouri ironweed	ASTERACEAE	herb	3
<i>Veronica arvensis</i> *	NI	VERO ARVE	corn speedwell	SCROPHULARIACEAE	herb	5
<i>Veronica peregrina</i>	FAC+	VERO PERS	necklace weed	SCROPHULARIACEAE	herb	5
<i>Vicia sativa</i> *	FACU	VICI SATI	common vetch	FABACEAE	herb	5
<i>Vicia villosa</i> *	no data	VICI VILL	vetch	FABACEAE	herb	19
<i>Viola sagittata</i>	FAC	VIOL SAGI	arrowleaf violet	VIOLACEAE	herb	24
<i>Vitis cinerea</i>	FACW	VITI CINE	grayback grape	VITACEAE	herb	23
<i>Vitis vulpina</i>	FAC+	VITI VULP	fox grape	VITACEAE	woody vine	3
<i>Vulpia octoflora</i>	FACU	VULP OCTO	sixweeks fescue	POACEAE	herb	21
<i>Wolffia brasiliensis</i>	OBL	WOLF BRAS	wolffia	LEMNACEAE	herb	11
<i>Xanthium strumarium</i>	FAC	XANT STRU	cocklebur	ASTERACEAE	herb	6

Species in bold type are tracked by the ANHC. (n = 12)

Species in red font are new additions to the master list for 2020. (n=7)

*/** = non-native/invasive species (99/496 = 20% of total)

+ = native species intentionally introduced to site (n = 7)

STRATA:

tree = ≥ 5 in dbh and ≥ 20 ft tall

sapling = 0.4 to < 5 in dbh and ≥ 20 ft. tall

shrub = usually 3 to 20 ft tall; multi-stemmed brushy shrubs, small trees, and saplings

woody vine = vines that are woody

herb = graminoids, forbs, ferns, fern allies, herbaceous vines, tree seedlings

SOURCE CODES:

1 = Chris Reid, site inventory, 17 August 2001 (west side)

2 = Bruce Shackleford, plot data

3 = Theo Witsell, site inventory, 1 August 2006

4 = Theo Witsell, Fall 2006 monitoring & inventory

5 = Theo Witsell, June 2007 monitoring & inventory

6 = Theo Witsell, October 2007 monitoring & inventory

7 = Theo Witsell, May/June 2008 monitoring & inventory

8 = Theo Witsell, September 2008 inventory

9 = Theo Witsell, November 2008 monitoring & inventory

10 = Theo Witsell, July 2009 monitoring & inventory

11 = Theo Witsell, October/November 2009 monitoring & inventory

12 = Theo Witsell, July 2010 monitoring & inventory

13 = Theo Witsell, October/November 2010 monitoring & inventory

14 = Theo Witsell, July 2011 monitoring & inventory

15 = Theo Witsell, November 2011 monitoring & inventory

16 = Theo Witsell, June 2012 monitoring & inventory

17 = Theo Witsell, November 2012 monitoring & inventory

18 = Bruce Shackleford & Seth Pickens, Spring 2013 inventory

19 = Theo Witsell, June 10 & 11 2013 monitoring & inventory

20 = Theo Witsell, November 2013 monitoring & inventory

21 = Theo Witsell, July 2014 monitoring & inventory (with Rubus identified by Dr. Johnnie Gentry, U of A)

22 = Theo Witsell, November 2014 monitoring & inventory

23 = Theo Witsell, June 2015 monitoring & inventory

24 = Theo Witsell, November 2015 monitoring & inventory

25 = Theo Witsell, 13 May 2016 site visit

26 = Theo Witsell, 19 May 2016 site visit

27 = Theo Witsell, 3 & 4 July 2016 monitoring & inventory

28 = Theo Witsell, 5 & 6 November 2016 monitoring & inventory

29 = Jeff Hickie, 2017 Growing Season

30 = Jeff Hickie, 2018 Growing Season

31 = Karen Willard & Jeff Hickie, 2019 Growing Season

32 = Karen Willard & Jeff Hickie, 2020 Growing Season

Appendix VI
Woolsey Wet Prairie
2020 Adaptive Management Activities

Woolsey Wet Prairie 2020 Adaptive Management Activities:

Date	Activity
February 5, 2020	2020 WWP Annual Management Meeting with City & Jacobs Personnel – ECO
February 21, 2020	2020 Prescribed Burn – OERI
March 25 & 26, 2020	Spot Spray yellow rocket, Queen Anne’s lace, curly doc, burdock, & motherwort w/ PastureGard HL – Jacobs
March 25 & 26, 2020	Spot spray tall fescue, winter bromes, & winter annuals w/ Section 2EC – Jacobs
April 2, 7, 9, & 14, 2020	Spot spray tall fescue, bromes, & winter annuals w/ Section 2EC – Jacobs
April 2, 7, 10, 15, 21, & 29, 2020	Spot spray yellow rocket, white sweet clover, burdock, curly doc, thistle, motherwort, & Queen Anne’s lace w/ PastureGard HL – Jacobs
May 1, 19, 20, & 29, 2020	Spot spray tall fescue, bromes, & winter annuals w/ Section 2EC – Jacobs
May 1, 5, 6, 11, & 18, 2020	Spot spray Callery pear, blackberry, winter creeper, bush honeysuckle, privet, honey locust, green ash, tree of Heaven, multiflora rose & persimmon w/ Remedy Ultra – Jacobs
May 7, 19, 20, 28, & 29, 2020	Spot spray yellow rocket, white sweet clover, burdock, curly doc, thistle, motherwort, & Queen Anne’s lace w/ PastureGard HL – Jacobs
June 2, 3, 5, 6, & 8, 2020	Karen Willard growing season invasive species and tracking species monitoring event
June 1, 2, 3, 4, 18, 22, 23, 24, & 29 2020	Spot Spray sericea lespedeza, Queen Anne’s lace, white sweet clover, & thistle w/ PastureGard HL – Jacobs
June 4, 5, 25, & 26, 2020	Spot spray Callery pear, blackberry, narrow-leaf cattail, willow, honey locust, green ash, multiflora rose & persimmon w/ Remedy Ultra – Jacobs
July 1, 2, 3, 8, 10, 14, 20, 21, 23, 27, & 28 2020	Spot Spray sericea lespedeza, Queen Anne’s lace, white sweet clover, & thistle w/ PastureGard HL – Jacobs
July 1, 6, 7, 20, 23, 28, & 30, 2020	Spot spray Johnsongrass w/ Outrider – Jacobs
July 20, 21, 23, 27, & 28 2020	Spot spray small carpetgrass w/ Section 2EC – Jacobs
July 6, 20, 21, & 23, 2020	Spot spray Callery pear, blackberry, narrow-leaf cattail, willow, honey locust, green ash, tree of Heaven, & persimmon w/ Remedy Ultra – Jacobs
August 4, 5, 24, & 26, 2020	Spot spray Callery pear, blackberry, narrow-leaf cattail, willow, honey locust, green ash, multiflora rose, & persimmon w/ Remedy Ultra – Jacobs
August 2020	Spot spray sericea lespedeza w/ Remedy Ultra – Jacobs
August 2020	Spot spray small carpetgrass w/ Section 2EC – Jacobs
August 25, 26, 27, 28, 29, 30, & 31 2020	Karen Willard post growing season invasive species and tracking species monitoring event
September 2020	Spot spray sericea lespedeza w/ Remedy Ultra – Jacobs
September 2020	Spot spray small carpetgrass w/ Section 2EC – Jacobs
October 2020	Spot spray black berry & Japanese honeysuckle w/ Remedy Ultra – Jacobs
October 2020	2021 Prescribed Burn Preparations Initiated – Jacobs
November 2020	2021 Prescribed Burn Preparations Completed – Jacobs
December 2020	Completion of 2020 WWP Annual Monitoring Report that includes 2021 Management Strategy – ECO

Appendix VII
2020 Woolsey Wet Prairie
Surplus Wetland Credit Ledger Report

CITY OF FAYETTEVILLE WOOLSEY WET PRAIRIE SURPLUS WETLAND CREDITS LEDGER
SERVICE AREA: City Jurisdictional Property Within Illinois River Watershed 8-digit Hydrologic Unit Code (HUC) Watershed (11110103)
Contact: Tim Nyander, City of Fayetteville Utilities Director
113 West Mountain Street
Fayetteville, Arkansas 72701
479-575-8390


Entry #	Entry Date	Available Credits	Withdrawn Credits for Impact Site	Debit (Impact) Site	Date of Credit Withdrawal	Debit Site Impacts	Debit Site COE Project/Permit No.	New Credit Balance
1	06/30/13	20.90 ¹	3.09 ²	Van Asche Drive Extension Terry Gulley, City of Fayetteville Transportation Services Director 479-444-3491	June 30, 2013	0.31 acres Palustrine Emergent Wetland	2012-00525	17.81
2	01/05/15	20.90	2.94 ³	Van Asche Drive Extension Terry Gulley, City of Fayetteville Transportation Services Director 479-444-3491	October 10, 2013	0.31 acres Palustrine Emergent Wetland	2012-00525-1	17.96
3	04/16/15	NA	NA	Correcting entry to change Van Asche Drive Extension contact to be Chris Brown City of Fayetteville City Engineer 479-575-8207	NA	NA	NA	NA
4	04/16/15	17.96	3.14	Clabber Creek Recreational Trail Chris Brown City of Fayetteville City Engineer 479-575-8207	April 10, 2015	0.192 acres Emergent Wetlands	2013-00322-1	14.82
5	12/16/20	14.82	2.50	Clabber Creek Recreational Trail Chris Brown City of Fayetteville City Engineer 479-575-8207	December 16, 2020	0.153 acres Emergent Wetlands	2006-00011470	12.32

¹As per September 30, 2013 Corps Correspondence approving use of Woolsey Wet Prairie 20.90 surplus wetland credits for impacts to wetlands caused by municipal projects within the Illinois River Watershed 8-digit Hydrologic Unit Code (HUC) watershed 11110103 (Standard Permit Modification No. 1997-14207-3) contained as Attachment #1 in January 5, 2015 City of Fayetteville, AR Woolsey Wet Prairie Surplus Wetland Credit Guidance.

²As per February 14, 2013 initial mitigation credit work sheets submitted to Corps contained as Attachment #2 in January 5, 2015 City of Fayetteville, AR Woolsey Wet Prairie Surplus Wetland Credit Guidance.

³Correcting entry for final required credits for mitigation as per October 10, 2013 Van Asche Drive Project Corps of Engineers Section 404 Permit No. 2012-00525-1 contained as Attachment #3 in January 5, 2015 City of Fayetteville, AR Woolsey Wet Prairie Surplus Wetland Credit Guidance.

[illegible]



REPLY TO
ATTENTION OF

DEPARTMENT OF THE ARMY
LITTLE ROCK DISTRICT, CORPS OF ENGINEERS
POST OFFICE BOX 867
LITTLE ROCK, ARKANSAS 72203-0867
www.swl.usace.army.mil/

Regulatory Division

STANDARD PERMIT MODIFICATION NO. 1997-14207-3

The Honorable Lioneld Jordan
Mayor of Fayetteville
113 West Mountain Street
Fayetteville, Arkansas 72701

Dear Mayor Jordan:

This letter is in response to the request by Environmental Consulting Operations, Inc. (ECO) to recalculate impacts incurred and mitigation required for Department of the Army (DA) Permit No. 1997-14207. The project site is located in the E ½ of section 14, T. 16 N., R. 31 W., in Fayetteville, Washington County, Arkansas.

Regulatory personnel have evaluated the original authorizations, existing site conditions, and current proposal. The impacts and mitigation credits were reviewed and recalculated based off of the original 2005 mitigation proposal, with these changes/additions:

1. On the Adverse Impacts calculation, corrected Duration value for Wastewater Treatment Plant and North Broyles Road from 0.2 to 2.0.
2. On the Adverse Impacts calculation, adjusted impact acreages based on information from ECO in June 2013.
3. On the Adverse Impacts calculation, adjusted Cumulative Impact value to account for lesser impact acreage.
4. On the Restoration and Enhancement calculation, removed Eastside Line Work as there were no impacts to restore.
5. On the Restoration and Enhancement calculation, changed the Net Improvement value for the buffer areas to 0.1, per ECO.
6. On the Restoration, Enhancement, and Creation calculations, separated herbaceous and forested as well as inside berm and outside berm to maintain consistency with original proposal.
7. On the Restoration, Enhancement, and Creation calculations, adjusted the Control value to "Covenant POA," except for Westside Line Work, which cannot be deed restricted.
8. On the Creation calculation, adjusted wetland created acreage to account for additional wetlands created.
9. On the Creation calculation, adjusted Vegetation values within the berms to 0.25, and used a value of 0.1 for areas outside the berms, per ECO.

As identified on the attached 2002 Charleston Method calculation sheets, 73.57 credits are

required to mitigate for impacts to aquatic resources for the City of Fayetteville's Wastewater Treatment Plant. We have calculated that the mitigation areas have generated 94.47 wetland credits. Therefore, the city will be able to use the excess 20.90 wetland credits to mitigate for wetland impacts generated by the City of Fayetteville within the Illinois River watershed, HUC 11110103.


The additional areas (referred to as South, West, and North Buffer) cannot generate buffer credit for this project. If you are interested in expanding the mitigation area into these areas, please submit a mitigation bank prospectus and we will evaluate these areas at that time.

This project and the Woolsey mitigation area present a unique situation in which we are considering new assessments of wetland impacts for a finalized project and recalculation of credits generated from a completed mitigation area. Please note that the Corps Regulatory Division does not intend to use this approach with other permit actions. It would not be feasible to make this a standard practice with the numerous issued permits, mitigation sites, and wetland banks finalized within the Little Rock District. The mitigation assessment credits for this 43-acre site will not be reconsidered in the future.

This letter becomes a part of and should be attached to your original permit.

If you have any questions, please contact Lisa Boyle, Project Manager, at (501) 324-5295 and refer to DA Permit No. 1997-14207-3.

Sincerely,



M. Elaine Edwards
Chief, Regulatory Division

Enclosures

CERTIFIED MAIL - RETURN RECEIPT REQUESTED

Copy Furnished:

Environmental Consulting Operations, Inc.

Mr. Bruce Shackleford, w/cy permit

Arkansas Department of Environmental Quality, w/cy dwgs

Proj Mgr, Beaver Lake PO, w/cy permit

Ch, Regulatory Enf, w/cy permit

Mr. Rocky Presley, w/cy permit